

Store  
HA O2823

Glasgow  
University Library



Store  
HA O2823



30114004659036

Glasgow University Library

GUL

~~1921/1998~~

12 FEB 2003

GUL 68.18



GENERAL BOARD OF HEALTH.

---

MEDICAL COUNCIL.

---

REPORT

OF THE

COMMITTEE FOR SCIENTIFIC INQUIRIES

IN RELATION TO

THE CHOLERA-EPIDEMIC OF 1854.

---

Presented to both Houses of Parliament by Command of Her Majesty.

---



LONDON:


PRINTED BY GEORGE E. EYRE AND WILLIAM SPOTTISWOODE,

PRINTERS TO THE QUEEN'S MOST EXCELLENT MAJESTY.

FOR HER MAJESTY'S STATIONERY OFFICE.

---

1855.



## GENERAL CONTENTS.

---

	Page
REPORT OF THE COMMITTEE FOR SCIENTIFIC INQUIRIES - - - - -	5

---

### SUPPLEMENT :—

No. I.—Letter from President of the Board of Health to Medical Practitioners, with Instructions and Forms for Returns of Cases of Diarrhoea and Cholera, and Memorandum thereon - -	68
No. II.—Statistical Tables and other Documents, illus- trating the Report of the Committee for Scientific Purposes - - - - -	87
No. III.—Summary Table of Houses, Population, and Mor- tality in the Golden Square Districts - -	119
No. IV.—Pathological Memorandum, circulated among Medical Officers of Public Institutions and other Members of the Medical Profession - -	127

---

## CONTENTS OF REPORT.

Duties and Objects of the Committee	-	-	-	Page 5
 <i>I.—Statistics.</i>				
Statistical Material employed	-	-	-	6
Progress and Fatality of the Epidemic	-	-	-	7
Estimated Number of Attacks	-	-	-	8
Local Differences of Cholera Mortality	-	-	-	11
Influence of Density of Population	-	-	-	12
Influence of Elevation	-	-	-	13
Mortality of Attacks of Diarrhoea and Cholera in its various Forms				16
Duration of Cases of Cholera and Diarrhoea	-	-	-	20
Influence of Age and Sex	-	-	-	22
Variations of Fatality during the progress of the Epidemic	-	-	-	22
Comparison of Mortality in the two Epidemics	-	-	-	23
 <i>II.—Ætiology.</i>				
A.—Atmospheric Causes :				
1. Mr. Glaisher's Meteorological Report	-	-	-	24
Estimate of its Results	-	-	-	31
2. Special Examinations of the Atmosphere by Dr. Thomson and Mr. Rainey	-	-	-	34
Estimate of their Results	-	-	-	36
3. Sanitary Inspection of Metropolitan Districts	-	-	-	37
B.—Water-supply of London :				
1. Dr. Thomson's Chemical Inquiry	-	-	-	39
2. Dr. Hassall's Microscopical Inquiry	-	-	-	42
Estimate of their Results	-	-	-	44
General Remarks on the Causation of Cholera	-	-	-	48
§ Supplementary Remarks on the Outbreak in Soho	-	-	-	49
 <i>III.—Practical Pathology.</i>				
Necessity for such Inquiry, and Materials obtained	-	-	-	53
1. Through what channel does the exterior Cause of Cholera affect the Human Body ?	-	-	-	54
2. Has the Disease a Period of Incubation ?	-	-	-	57
3. Is it communicable from Person to Person ?	-	-	-	58
4. What relation does the Gastro-intestinal Flux bear to the other Phenomena of Cholera ?	-	-	-	59
5. What is the Pathology of the Consecutive Fever ?	-	-	-	61
6. What are the import, tendency, and characteristics of the epidemic Diarrhoea ?	-	-	-	62
7. Examinations of Blood, &c. &c.	-	-	-	65
 <hr style="width: 20%; margin: 10px auto;"/>				
SUGGESTIONS FOR FUTURE INQUIRIES	-	-	-	65



## R E P O R T.

---

*To the Right Hon. the President of the General Board of Health.*

SIR,

London, July 14th, 1855.

Soon after your having constituted a Medical Council to advise the Board of Health on matters relative to the then prevailing epidemic of cholera, this Council thought it convenient for the despatch of business that subjects referred to them should be distributed among certain committees of their number; and on this plan we, the undersigned, were requested to become a committee for the scientific purposes of the Council.

Two special duties accordingly devolved on us: *first*, to suggest the institution of particular scientific inquiries which we thought likely, by bettering medical knowledge of the disease, to strengthen the public resources for its prevention and cure; *secondly*, to review, as laid before us, the various fruits of this investigation, and to submit to you our judgment of its results.

While proceeding to report to you on our fulfilment of this task, we beg to acknowledge very gratefully the kindness and confidence with which you have honoured us, in instituting all the inquiries we ventured to suggest, and in procuring from other departments of Government whatever assistance could conduce to our success.

At the same time, we ask indulgence for much that is unavoidably imperfect in our work. To do such things fitly, all needful organization must be prior to the emergency; it would be vain to expect that observations begun in the crisis of an epidemic should have that completeness which science requires, and which only deliberate preparation can ensure. We need hardly remind you that the re-constitution of the General Board, under which you became its President, was effected only on the 12th of last August; that the first meeting of the Medical Council was only on the 6th of September; and that we were appointed a committee two days later—already in the tenth week of a pestilence which some days previously had

attained its utmost extension, and was now in process of decline.

If, however, we cannot speak with unmixed satisfaction of the materials which are before us, we can at least point to their nature and extent as in the highest degree encouraging to future exertions, more extended, more systematic, and more continuous.

---

Our principal aims, and the methods by which their attainment was sought, have been as follows:—

I. with a view to the DESCRIPTIVE HISTORY OF CHOLERA, we have examined the larger *statistics of this invasion*; as to the places wherein the disease chiefly prevailed; as to the influence of age, sex, and employment in favouring its attack; and as to its own pathological stages and periods;—

II. in the hope to gain more precise knowledge of the CAUSES OF THE DISEASE, we thought it of primary importance that the *air and water of the metropolis* during the epidemic period should be studiously observed, and that special inquiries should, as far as possible, be made into the state of these universal influences in districts actually infected with cholera;—and

III. with the object of increasing for our profession the present insufficient resources of MEDICAL TREATMENT, we have endeavoured to procure comparative *records of various therapeutical experience*, successful or unsuccessful, and have invited from persons versed in such inquiry an elucidation of those questions in the *practical pathology of cholera* which appeared to us most urgent for solution.

---

## FIRST SECTION.

---

### *Statistics.*

---

OF statistical material, the following has been before us:

1. We have derived from the General Register Office a *list of all deaths* registered during the epidemic period as caused by cholera or diarrhœa, with particulars as to the

age, sex, and residence of each sufferer, the date of death and, in many cases, the duration of fatal illness.

2. As the *returns of certain circular forms* (A. and B.) distributed by the Board, we have received from more than 300 medical practitioners, whose names we append to our report, an immense mass of detailed information; and to this we cannot advert without offering our tribute of respect to the public spirit which led so many members of our profession, solely for the general good, to incur an additional task during days of overwhelming occupation. These returns relate to 4,271 cases of cholera and 20,301 of diarrhœa. Besides particulars hereafter to be mentioned in respect to the treatment of the disease, there is contained in them pathological information as to its stages and periods, as to the universality or relative frequency of certain symptoms, as to the prospects of death or recovery at each step of its progress.

From the above sources we gather the following facts of the late epidemic visitation—the third which London has suffered from Asiatic cholera.

### (i.) *Progress and Fatality of the Epidemic.*

The summer of 1853 witnessed its commencement.\* At that time, as the temperature rose in July, diarrhœa, as well as the common form of cholera, became fatal here; and a few deaths from cholera in the Asiatic form were registered in August in the low districts by the side of the river. Several deaths by the disease occurred in September; and in October, while the temperature fell, and diarrhœa decreased from 723 in August to 283 in October, the cholera spread and became more fatal, so that the deaths by cholera were 335 in October, and 288 in November. On the last day of October 25 persons died of the disease; but subsequently the epidemic subsided, so that towards the close of November the deaths did not exceed four daily; in the month of December the deaths were 43; in January (1854) one death only happened at intervals on each of seven days; one death was recorded on the 10th of February, one on the 25th, and one on the 26th; no death by cholera occurred in March; only four in April, four in May, and three in June.

---

\* Earlier in the year, the disease had prevailed in the ports of the Baltic and on the shores of the German Ocean.

Diarrhœa, however, although it had declined, never ceased; no day in the six first months of the year was without a death, and on some days as many as 8, 10, and 12 deaths by this form of disease were recorded; but it exhibited no disposition to increase. Yet, warned by the course of the former outbreaks, fears were entertained of the more formidable secondary visitation; which were unfortunately justified by the event, for in the eleven days after July 4th, ten persons died of cholera; on the 16th, four died of the disease; and the deaths ran up towards the end of the month until 53, 41, and 44 died on the last three days; on August 1st, 72 persons died of cholera; which continually spreading and multiplying in its course, the deaths on the last day of August were 211. The deaths in July had been 371 by diarrhœa, 308 by cholera; in the month of August the deaths by diarrhœa amounted to 1,022, by cholera to 3,513. No abatement was observed, but the epidemic raged more violently, and the deaths by cholera alone on the 1st of September were 389; on the 2nd, 459; on the 3rd, 329; or 1,177 in the three days, beside 126 deaths by diarrhœa. The eruption had now reached its culminating point; but it subsided slowly, for 4,371 lives were destroyed by cholera and diarrhœa in the first fourteen days of September, while in the month the deaths by cholera were 6,084, by diarrhœa 990. The epidemic declined rapidly in October, and the deaths fell from 70 by cholera on the 1st day, to 3 on the 30th day, making 823 in the aggregate, besides 426 by diarrhœa. In November the deaths by cholera were 52; in December 5, namely, one on the 2nd day, one on the 5th, one on the 6th, and two on the 22nd day. The deaths by diarrhœa in the two months were 175 and 113. The plague was stayed, but it had destroyed in one or other of its forms seventeen thousand people. Such is a rapid sketch of this remarkable epidemic, which can be traced in the annexed Tables, through its fatal course, day by day, from July 1st, 1853, to the end of the year 1854. (See Table I., Second Series.)

(ii.) *Estimated Number of Attacks.*

The deaths by the disease were all registered; and the names, ages, occupations, death-dates, and death-places of



its victims are all recorded in the books of the General Register Office. The list of the killed is therefore complete; but what was the number of the wounded? What number of the people was attacked by cholera, and what number suffered and recovered from diarrhœa? No complete return of the cases exists; but the forms of return (A. and B.) which were issued by the Board of Health, and filled up in a manner so creditable to the medical practitioners of London, enable us to form an estimate, which cannot differ much from the actual facts.

Thus, the medical returns show that of 3,188 recorded cases of cholera, 1,467, or 46 per cent., terminated fatally; so it may be inferred from this proportion that, as 11,661 persons died, about 25,000 were attacked by cholera. The mortality of the cases in the hospitals was 51 per cent. ( $\frac{763}{1502}$ ); the mortality of cases that were treated at home was 42 per cent. ( $\frac{704}{1686}$ ); and some deviation further still from the average may have actually occurred in the whole population, but the estimate is true within certain limits. Again, 5,271 cases of diarrhœa were recorded in detail, distinguishing the ages, and 87 were fatal, or the mortality among persons actually attacked by diarrhœa was at the rate of .0165, about 1.65 per cent. And 17,351 cases, 109 deaths by diarrhœa (.00628) may be noted, if we count the additional cases that are returned in numbers without being separately entered. The deaths in London from diarrhœa were 6,258, so it is evident that only a small proportion of these fatal cases—occurring chiefly in children and old people—attracted the attention of the medical observers. But we are already justified in inferring that, as 6,258 died, some *hundreds of thousands* of the population were attacked by the disease.

The medical returns show the power of recovery from an attack of cholera at different ages; so that 35 deaths at the age 15 to 25 imply that 100 persons of the age have been attacked; at the age 45 to 55, 50 deaths imply 100 attacks; at the advanced age, 75 to 85, 71 deaths by cholera imply 100 attacks. So it is with diarrhœa. And if the proportional numbers of deaths to cases in the medical returns are applied to the total deaths that were registered at the corresponding ages, the result gives the number that were attacked at all ages by cholera as 24,917 persons; by diarrhœa of some severity as 329,778; by diarrhœa of so slight a nature as to be only

brought casually under medical observation, about 519,487; making 874,182 persons in the aggregate who were touched by the epidemic, while 1,642,866 persons escaped unscathed. The estimate of the slight cases is based upon imperfect observations; and we set it down here as a mere indication of the wide influence of the epidemic. (*See First Series, Table XII.; Third Series, Table VIII.*)

The aggregate number of deaths by diarrhœa and by cholera in the medical returns was 1,576, of cases 20,648; and if the estimate is taken on these proportions, it will follow that the 17,919 deaths by the two forms of disease imply less than 235,000 cases; but it is evident on the face of the returns that the slighter cases of diarrhœa were in many instances unrecorded, and consequently the severer forms of cholera were returned by the medical observers in a higher proportion than they actually occurred; we have, therefore, made the above estimate on the separate returns. There can be no doubt, however, that, as is subsequently shown, the great majority of the registered deaths from diarrhœa were deaths from cholera in one of its modified forms; and that both these deaths and all those registered as deaths from cholera, should be taken as representing the mortality of one epidemic disease, including cholera and diarrhœa. To arrive at a correct estimate, however, of what the rate of mortality of the entire disease (including cholera and diarrhœa) is, a large addition must be made to the above number of cases for cases of diarrhœa omitted.

London, in the middle of the year 1854, contained about 2,517,048 people; whom we may conceive to be distributed in equal groups of 10,000. The epidemic diffused itself over each of the 251 myriads; and if *one myriad* is taken to represent the average danger and suffering of the whole, it appears that 71 died; namely, 46 by cholera, 25 by diarrhœa; 99 having probably been attacked by cholera, 1,310 by diarrhœa of some severity, and a still larger number (perhaps 2,064) by slighter forms of disorder. It may, indeed be assumed that in one period or other of the epidemic every individual fell more or less under its influence; while to some, however, who tasted the poison, it was only the cause of a temporary derangement, to others who drank of the chalice to the dregs it was the bitterness of death.

In the years 1840–1 the deaths from diarrhœa in London

amounted to 452 and 465; in the four years 1842-5 the deaths fluctuated from 704 to 834; and since the year 1846 this disease has, in the years when cholera was not epidemic, been the cause of death to 2,000 or more of the inhabitants, and must have attacked 100,000 of them annually, or 75,000 more than it did in 1840-1. Summer cholera, within the same period, also became more prevalent; 60 deaths were ascribed to that cause in 1840, and 162 in 1852. A certain number of the deaths from diarrhœa and cholera in the period of the epidemic are therefore referable to these diseases in the old form, and a certain number to the form acquired since the hot summer of 1846. No inconvenience will, however, arise in the subsequent inquiries if the rate of mortality is calculated on the population in 1854, and on the deaths from cholera and diarrhœa in the period from July 1st, 1853, to December 31st, 1854.

During the epidemic many persons labouring under chronic diseases are attacked by diarrhœa and cholera, which prove fatal in a certain number of instances; and in the abstracts that we have had made from the public registers these cases are counted, although the original primary disease was of a fatal nature. Our numbers, therefore, will exceed the numbers referred to diarrhœa in the Registration Tables.

### (iii.) *Local Differences of Cholera Mortality.*

The rate of mortality all over London is represented by the deaths of 71 persons in a myriad people in average conditions of exposure. We cannot investigate the mortality in each of the 251 myriads of which London is composed, but we have the means of determining in the usual way the mortality in each of the 36 registration districts, containing populations varying from one (Hampstead, 11,986) to sixteen (Pancras, 166,956) myriads. Here the hospitals interfere to some extent; but a correction has been made by distributing the 800 deaths in the hospitals over the several districts, in the proportion of the deaths that occurred out of the hospitals in each district, so that the disturbance from this cause cannot be considerable. A similar inquiry has been undertaken, for the sake of comparison in each of the 135 sub-districts into which the 36 districts are subdivided; and some cor-

rection has been made for the deaths from cholera in the workhouses, which in certain districts were used as cholera hospitals. (*See Second Series, Tables III., IV., VI.; Third Series, Table I.*)

The districts, arranged in the order of the rate of mortality from cholera, display a regular series of numbers expressive of that rate, ranging from 6, 10, and 11 at one extreme, to 142, 165, and 179 at the other extreme; so each myriad of the people in the different localities of this great city suffered differently, and the observations present us with these enormous disparities in the sufferings of the thirty-six large congregated masses. The range in the rate of mortality by diarrhœa is much less considerable; it extends from 11, 12, 17, in some districts, to 39, 53, and 54, in a myriad inhabitants, in other districts.

#### (iv.) *Influence of Density of Population.*

The population of London stands on 78,029 acres of ground, so that in 1851 there were on an average 30 persons to an acre. In Lewisham there were 2 persons, in Wandsworth 4, and in Hampstead 5 persons to an acre; the mortality to 10,000 from cholera in these districts was at the rate of 22, 85, and 12 in 1853-4; and 30, 100, and 8 in 1849 in these open districts. In the three densest districts there were 246, 256, and 290 persons to an acre; the mortality from cholera was at the rate of 10 in St. Luke, of 22 in the Strand, and of 23 in the East London City district, in 1853-4; of 34, 35, and 45 in the same districts in 1849. The mean mortality by cholera is, in the two epidemics, at the rate of 43 in the three most open districts, 28 in the three most dense districts.

Again, in the nine districts of Lewisham, Wandsworth, Hampstead, Camberwell, Hackney, Kensington, Poplar, Greenwich, and Rotherhithe, the mean density of the population ranges from 2 to 21 persons on an acre; the mean mortality in the two epidemics was at the rate of 65 by cholera, by diarrhœa 22, in 10,000.

And in the nine densest districts, Whitechapel, St. George in the East, the West London City, St. Giles, St. James, Holborn, St. Luke, the Strand, and the East London City, where there were on an average from 196 to 290 persons on an acre, the mean mortality in the two epidemics was, by cholera 41, by diarrhœa 19, in 10,000.



The mortality by the two forms of disease was 85 in the nine open to 58 in the nine dense districts.

The mean mortality by cholera and diarrhœa, in the 18 most open districts (40 in 10,000), is nearly the same as (42 in 10,000) the mortality in the 18 most dense districts.

If the 135 sub-districts are arranged in the order of their density, the result is similar; the fatality of the epidemic being highest, however, in the districts of an intermediate degree of density. The cholera matter was evidently diffused over every sub-district of London, but it does not appear that the great differences in the density of the habitations of the people exercised any decisive influence on the intensity of its operation, appreciable, at least, by this method of investigation. Its effect was, perhaps, masked by other more potent agencies. This is the more remarkable as the fatality of large classes of disease has been found to increase in a given ratio to the density of the population; and it may be inferred that cholera is not, like some such cases, communicated by the breath from person to person. (*See Third Series, Tables I., II., III.*)

#### (v.) *Influence of Elevation.*

The population of London is distributed over the low ground on both sides of the Thames, and over a great number of elevations and depressions, which ascend from the south bank of the river up to Blackheath and Norwood, and from the north bank up to Highgate and to Hampstead. The four lowest districts, Newington, Rotherhithe, St. George Southwark, and Bermondsey, are on or below the level of the Thames at high water; the mortality by cholera to 10,000 in these districts was at the rate of 112, 165, 121, and 179 in the last, and 144, 205, 164, and 161 in the previous epidemic.

Hampstead, Islington, Marylebone, and St. Pancras are at average elevations of 350, 94, 87, and 73 feet above the Thames, and the mortality by cholera in these highest districts was at the rate of 12, 11, 17, and 10 in the last, of 8, 22, 17, and 22, to 10,000, in the former epidemic.

The mean mortality by cholera to 10,000 in the two epidemics was at the rate of 156 in the four lowest districts, 15 in the four highest districts. The mortality by *cholera and diarrhœa* to a myriad of population was 189 on the low districts, 34 on the high districts.

If the thirty-six DISTRICTS of London are arranged in the order of their elevation above the high-water mark of the Thames, the mortality by cholera is found not to be invariably in each district inversely as the elevation; but by taking groups of districts together in the two epidemics, a nearly regular series is obtained: thus, the mean mortality by cholera was, to every myriad, 156 in the districts on or below the level of the high-water mark, 91 in the districts of 3 and under 20 feet of elevation, 44 in the districts at 20—40 feet, 36 in the districts at 40—60 feet, 23 at 60—80 feet, 17 at 80—100 feet, and 10 at 350 feet of elevation. (*See Third Series, Table I.*)

The mortality by diarrhœa was at the rate of 33 on the lowest ground, 26 on the second terrace, 19 on the third terrace, 18 on each of the three higher terraces (20—100 feet), and 10 on the highest terrace. Thus, the mortality by diarrhœa varies less than the mortality by cholera at different elevations. Upon the two highest terraces the diarrhœa is as fatal as the cholera; upon the lowest ground the cholera is four times as fatal as the diarrhœa.

The relation between the elevation of the dwelling-ground and the intensity of the epidemic is seen in the annexed Tables of the Third Series.

(Table IV.) Of the 135 SUB-DISTRICTS, arranged in regular order from the highest to the lowest in London;

(Table V.) Of these sub-districts grouped together in fourteen terraces;

(Table VI.) And of the same sub-districts grouped in six terraces of elevation. The form is the same as the corresponding Tables in the Registrar-General's weekly tables; but the mortality is deduced from the deaths by cholera and by diarrhœa in the 18 months, July 1st, 1853, to the end of 1854; and certain corrections are made for the increase of population and for the disturbance that the deaths in hospitals and workhouses occasioned. The results in the Table V. and Table VI. present a near approximation to the true rates of mortality in the 135 sub-districts; and though differing in details, are of the same character as the results that have been deduced above by grouping the 36 districts. (Table I.)

The mortality from cholera (1853½—4) was at the rate of 13 to a myriad in the highest, 137 to a myriad in the lowest sub-districts (Table V.); the mortality in the same sub-districts from diarrhœa was 21 in the highest,

34 in the lowest ; and it will be recollected that *one* death from cholera represents about *two* (2·2) cases of cholera, while *one* death from diarrhœa represents about *sixty* cases of diarrhœa of some severity ; consequently the cases of diarrhœa and cholera together must at these rates have been about 1,288 to a myriad in the higher regions, and 1,741 in the lower regions of London. But if it be assumed, as is not impossible, that the cases of diarrhœa and cholera were less fatal on the elevated sub-districts than the cases on the low grounds, then the proportion of persons attacked in the respective regions would differ much less considerably : for, if 1 in 35 cases of diarrhœa was fatal in the lower regions, and 1 in 70 was fatal in the higher regions, the proportional number of persons that were attacked by diarrhœa or cholera in each must have been about 1,490 in a myriad of the population ; and taking intermediate proportions, a similar result is obtained for the regions at intermediate elevations.

The distribution of choleraic attacks (though in widely different degrees of frequency, and perhaps also of severity) throughout the whole metropolitan area, seems to establish that the *cholera-leaven*, be it what it may, was scarcely less diffused in the districts that suffered the lowest mortality, than it was in the districts where the disease was tenfold more fatal.

But while the presence of this *leaven* seems to have been universal throughout the districts of the metropolis, the consequences excited by its presence have greatly varied in different localities : and independently of any hypothesis, it may now be stated as the experience of two epidemics in London, that such local varieties of effect, grouped into masses for comparison, have been more nearly inverse to the elevation of soil in the affected districts than proportionate to any other general influence that we could measure.\* Thus, approaching London

\* The following formula is from the Report to the Registrar-General on the Cholera of 1848-9, p. lxiii. :—Let  $e$  be any elevation within the observed limits, 0 to 350 feet ;  $c$  be the rate of mortality from cholera at that elevation ; also let  $e'$  be any higher elevation, and  $c'$  the mortality at that higher elevation. Then, if the mortality from cholera is inversely as the elevation, we shall have the proportion,—

$$(1.) e : e' :: c' : c = \frac{e'}{e}, c'.$$

By adding a constant element,  $a$ , the velocity at which the mortality increases can be retarded to any extent. The equation then assumes the form,—

$$(2.) \frac{e' + a}{e + a} \cdot c' = c, \text{ or } (3.) c' = \frac{e + a}{e' + a} \cdot c.$$

along the roads from the surrounding country, and descending through the successive regions, succeeding each other in circles, down to the waters of the polluted Thames, we see, in the epidemic, the people fall upon the right hand and upon the left in numbers that increase on every circle, and express arithmetically the growing force of those physical influences, on which the poison of cholera apparently depends for its powers of existence or of development.

*The annexed diagram* exhibits to the eye the relative intensity of the cholera in the sub-districts of various elevations; it also exhibits a regular curve, with which the series observed closely agrees, except in the part which includes the observations in the Berwick Street and Golden Square sub-districts.

(vi.) *Mortality of Attacks of Diarrhœa and Cholera in its various Forms.*

The medical returns already adverted to, contained more or less complete information of the kind indicated by the Forms A, B, which were drawn up by us, in the midst of the epidemic. Those returns came to hand at distant intervals from the London Hospitals, from private practitioners, and from country districts; and the various sets of returns have been made available, as far as it was practicable, for the various branches of this inquiry; thus 24,572 cases, namely, 4,271 of cholera, and 20,301 of diarrhœa, were brought under our observation. The mortality by cholera in the several sets of English returns, ranged from 41 to 51 deaths on every 100 persons attacked; the mortality from diarrhœa ranged from .6 to 1.8 on every 100 attacked, (or without decimals, from 6 to 18 on every 1,000), as is shown in the annexed Table :

The value of  $a$  in general terms is,—

$$(3.) \ a = \frac{e' \ c' - e \ c}{c - c'}, \text{ and it was taken at 13.}$$

$e + a$  and  $e' + a$  represent the abscissas,  $c$  and  $c'$  the ordinates of the curve. The central perpendicular line in the diagram corresponds to the abscissas, the horizontal lines to the ordinates of the curve, which was calculated from the mean mortality (145) on the lowest ground ( $c = -1$ ), one foot below Trinity high-water mark. The equation was, therefore,—

$$\frac{a - 1}{a + e'} \cdot c = \frac{13 - 1}{13 + e'} \times 145 = c' = \frac{12 \times 145}{13 + e'} = \frac{1740}{13 + e'};$$

the  $e'$  being made the variable.



TABLE (C.) of MORTALITY of attacks of CHOLERA in several Groups of Returns.

—	Number of Cases.	Deaths.	Mortality. Deaths to 100 Cases.
(1.) Cases from all sources -	4,271	1,948	45·6
(2.) Cases in London Hospitals -	1,502	763	50·9
(3.) Cases in private practice, } London - - - }	1,686	704	41·8
(4.) Total of (3.) & (4.), or of } the London Cases - - }	3,188	1,467	46·0
(5.) Cases in the provincial towns } and districts - - }	586	238	40·6
(6.) Some cases in Scotland -	497	243	48·9
(7.) A selection of all cases in } which the connexion of } collapse Cases, or Cases } without collapse, with con- } secutive fever, could be } traced - - - }	3,596	1,767	49·1
(8.) Of the above (7.) were cases } of collapse* - - - }	2,431	1,627	66·9
(9.) Of the above (7.) were cases } without absolute collapse }	1,165	140	12·0
(10.) All cases in which age was } specified - - - }	3,611	1,749	48·4

TABLE (D.) of MORTALITY of Cases of CONSECUTIVE FEVER following CHOLERA (included in No. 7 of Table of Cholera Mortality).

—	Number of Cases.	Deaths.	Mortality. Deaths to 100 Cases.
All Cases of consecutive fever -	874	249	28·5
Of the above were :—			
(1.) Cases following collapse -	633	220	35·0
(2.) Cases following cholera } without collapse - - }	241	29	12·0

\* The numbers are taken from the Tables IV. and V., First Series.

TABLE (E.) of MORTALITY of Attacks of DIARRHŒA, Simple or Choleraic.

	Number of Cases.	Deaths.	Mortality. Deaths to 100 Cases.
(1.) Cases from all sources -	20,301	156	·77
(2.) Cases treated in London } hospitals - - - }	688	8	1·16
(3.) Cases not treated in London } hospitals - - - }	16,772	101	·60
(4.) Sum of cases in London -	17,460	109	·62
(5.) Cases in the country towns } and districts where cho- } lera was epidemic - - }	2,176	40	1·84
(6.) Some cases in Scotland -	665	7	1·05
(7.) Cases in which age was dis- } tinguished - - - }	5,271	87	1·65

From another extended series of returns that were procured through the Poor Law Board of Scotland, it appears that the mortality in that country among persons attacked by cholera, was 47·5 in 100; for out of 14,430 cases, 6,848 terminated fatally. This result agrees closely with the general result of the English cases, of which 45·2 in 100 were fatal. (*See First Series, Table XIII.*)

If we assume for a moment that all the cases commence as diarrhœa, we may infer that of 142,351 persons attacked of all ages, a certain number, say 132,351, suffer from diarrhœa of some severity, and that after the lapse of a certain number of days 2,512 die, 129,839 recover: but 10,000 enter the stage of cholera, and may then be divided into two great classes: 3,240 of the *first class*, who do not fall into collapse; 309 of their number dying, 2,261 recovering, while 670 pass into consecutive fever, 81 to die in, 589 to recover from, that stage; and 6,760 of the *second class* who do fall into collapse, 3,913 dying in that stage, from which, however, 1,087 recover straightway, leaving 1,760 who pass into consecutive fever, in which 612 die, and from which 1,148 recover: thus of the 10,000 that we follow in their perilous journey, 6,760 fall into collapse, 2,430 fall into the fever; of the 6,760 who fall into collapse, 4,525 die either in that stage or ultimately

in the fever stage: of the 3,240 who do not fall into collapse, as many as 390 die: of the 2,430 who pass into the fever 693 die.\*

If a single patient is regarded, these numbers will assist in prognosis, as they express the degrees of danger to which he is exposed; thus upon being attacked by unequivocal cholera, the probability is slightly in favour of recovery (.5085); but the chance is two to one that he will fall into collapse, and then it is two to one that he will not recover; if, however, he survive this stage and pass into the fever, it is more than two to one that he will recover. It is seven to one that the patient who does not fall into collapse will recover, and if he pass into the fever he has such an advantage that it is seven to one in favour of his recovery, while, as is before stated, it is only two to one in favour of the recovery from the fever of a patient who has been in collapse.

Such, it must be recollected, is not the natural course of these two forms of disease, but their course under the systems of treatment now in practice in England: and what is the result of the English practice?

The great object of the physician or surgeon in every stage of the malady is to save the patient's life, and with this object in view, to prevent the diarrhœa from passing into cholera, the cholera from falling into collapse. Now it has been seen that of 142,351 persons attacked, about 129,839 recover, while only 10,000 of the cases become cholera, 6,760 fall into collapse.

The question, how far these proportions have been affected, for better or for worse, by various methods of treatment, empirical or rational, has been under the con-

\* The following are the data from which the above results have been deduced:—

The number of severe cases of diarrhœa, in London, 329,778 (estimated); deaths by diarrhœa, 6,258; cases of cholera, 24,917 (estimated); cases of cholera in which the several stages were distinguished in the medical returns, 3,596, of which 1,767 terminated fatally.—(See First Series, Table IV.; Third Series, Table VIII.)

1,165 of the 3,596 cases did not go into collapse; but 111 died, and 813 recovered from this stage.

2,431 cases went into collapse, from which 391 recovered without proceeding further, and in which 1,407 died.

874 of the 3,596 cases passed into the stage of consecutive fever; and as it was observed that of 808 such cases, 223 had been preceded by cholera without collapse, 585 by cholera with collapse, it was inferred that of the 874 cases, 241 had not been preceded, that 633 had been preceded, by collapse.

249 of the cases of consecutive fever terminated fatally; and as it was observed that of 223 cases, 26 had not been preceded, that 199 had been preceded, by collapse, it was inferred that these two latter numbers should be raised to 29 and 220.

In a few rare instances the consecutive fever, it was stated, passed into other forms of disease, which are not brought into account in this illustration of the course of a complicated disease.—(See First Series, Tables IV. and V.)

sideration of some of our colleagues, acting as a special committee in this branch of the subject. We, therefore, only remark on one difficulty of the investigation which has been obvious in the materials we have analysed, and has often misled the public into false appreciation of alleged methods of cure. The choleraic pestilence varies in the severity of its individual attacks, from the degree of a trifling indisposition to that of a most deadly and intractable disease. We have seen that in one form it is fatal to 6, in another to 669 in every 1,000; and, therefore, to avoid great mistakes, any alleged specific requires that its effects should be investigated with the greatest care, through extended and, above all, trustworthy observations. When persons not accustomed to accurate investigations attempt to compare together the results of various treatment, as tested by death or recovery, they are seldom sufficiently on their guard against the immense fallacy of leaving unexpressed the *degrees of disease* against which this or that medicine has prevailed.

(vii.) *Duration of Cases of Cholera and Diarrhœa.*

If the first object of medical treatment is to save the patient's life, the second is to shorten his sufferings, and to accelerate his restoration to health. Now the duration of 3,600 cases of cholera, dating not from the commencement of the precedent diarrhœa, but from the first characteristic symptoms of cholera, was by the medical returns 5·9 (nearly six) days: the duration of the 1,744 fatal cases was 2·68 days (more than sixty-four hours); while the duration of the 1,856 cases of recovery was 9·06 days. We may, therefore, inquire of new methods of treatment, do they shorten the periods of the disease, as well as do they diminish the mortality more or less than the present system of practice?

The duration of 9,590 fatal cases of cholera was returned in the registers of deaths, and was on an average 2·39 days; a result differing little from that above, which is deduced from the medical returns.

If the series of changing phenomena in an attack of cholera are viewed collectively, with reference only to the time, dating from the commencement, we find, by a simple construction (*see* Third Series, Table IX.) that of 3,600 cases 816 terminate in one day (twenty-four hours), namely, 82 in recovery, 734 in death; leaving 2,784 cases,



of which 590 terminate in the second day, 162 in recovery, 428 in death; and so on to the 34th day as may be seen in the Table. The same construction shows that the average chance of recovery increases every hour, every day, from the commencement of the attack, at first rapidly, and then slowly: so that while it is only 1,856 to 1,744 in favour of recovery at the onset, it is 1,744 to 1,010 in favour of recovery if the patient survive twenty-four hours; the patient alive at the end of the second day, though still ill, has the chances of 1,612 to 582 (almost three to one) in favour of his ultimate recovery; at the end of ten days his chance of recovery is 636 to 109 or six to one: so that to gain time, is one great aim in the treatment of this disease, which destroys, in the first twenty-four hours after the manifest characteristic symptoms, *one* in every five that it attacks.

The probability that a patient suffering from an attack of average severity will die or recover in the attack is shown in Table XI. of the Third Series. In the doctrine of probabilities, certainty, it will be recollected, is expressed by *unity* (1), and the various degrees of probability are expressed by fractions, so that if an event can only terminate in one of two ways, the sum of the two fractions that measure the respective probabilities is 1·0; as by hypothesis the case must certainly terminate in one of the two ways. Thus, by the Table, the probability that a cholera patient will die of the attack at first is nearly  $\frac{48}{100}$  or ·48; at the end of two days the probability is reduced to ·27, at the end of the seventh day to ·16, at the end of nineteen days to ·07. The probability of ultimate recovery increases as the probability of dying decreases, and is at the corresponding dates ·52, and ·73, and ·84, and ·93.

The Table IX. (Third Series) shows at a glance the probability that a patient suffering from an attack of average severity will recover *on*, *before*, or *after* any day of the disease; and will serve, therefore, either to guide an insurance office in insuring the life of a patient or to direct the physician in determining the effects of various systems of treatment. We shall not pursue this branch of the inquiry further into details, which can only be made clear by the aid of mathematical symbols. But we may here remark, as the same method is applicable to all diseases, that medicine is a science of probabilities, having observation for its basis; and depending, therefore, for its success on the application of the same methods of analysis as are

applied in the other sciences to facts observed on some such extended system as we have endeavoured to illustrate in conducting the inquiry into the laws to which an epidemic disease is subject.

The average duration of 5,271 cases of *diarrhœa*, chiefly of recoveries in the medical returns, was five days, while the duration of 4,150 cases in the registers of deaths was 13 days, these being all severe and more protracted diseases, occurring chiefly in young and in very old people.

The duration of cases of *cholera* is by the medical returns nearly 6 days, of cases of *diarrhœa* nearly 5 days; so that the two forms differ little in duration, and the 329,778 cases of severe *diarrhœa*, which, if we take the previous estimate, occurred in London, imply about 1,648,890 days of sickness, while the 24,917 cases of *cholera* imply 149,502 days of sickness.

#### (viii.) *Influence of Age and Sex.*

The influence of age on the course of *cholera* is striking; thus in infancy and advanced age the form of *diarrhœa* is relatively and absolutely more frequent than it is in the middle period of life, as at that age the spasms and the collapse are more evident than they are when the muscular system is feeble.

The danger of an attack of *cholera* varies with age; thus at the age of 15–25, out of 100 persons attacked 34·9 die; at the age of 25–35 the deaths to 100 cases are 35·4; at 65–75 the deaths to 100 cases are 58·2. The mortality of cases of *diarrhœa* also varies at different ages.

In another series of tables the mortality that *cholera* caused in the population at various ages is shown from the facts of the two epidemics. (*See First Series, Table XII.; Second Series, Table X.*)

The mortality among the male population was at the rate of 47, among the female population at the rate of 45 in 10,000.

#### (ix.) *Variations of Fatality during the progress of the Epidemic.*

The fatality of the cases of *cholera* diminished in the progress of the epidemic, so that the disease apparently assumed towards its close some of the characters of *diarrhœa*; as is shown in the annexed tabular statement of cases returned in London:—

Period of the Epidemic.		Cases.	Deaths.	Deaths to 100 Cases.
Fortnights.	Dates.			
	Total Cases - -	3,188	1,467	46
1	July 2—15 - -	6	2	33
2	July 16—29 - -	41	24	59
3	July 30—Aug. 12 -	272	141	52
4	Aug. 13—26 - -	420	181	43
5	Aug. 27—Sept. 9 -	980	509	52
6	Sept. 10—23 - -	921	393	43
7	Sept. 24—Oct. 7 -	377	151	40
8	Oct. 8—21 - -	118	54	46
9	Oct. 22—Nov. 2 -	39	9	23
10	Nov. 3—16 - -	14	3	21

If periods of four weeks are taken, it will be found that the fatality of the cases in the returns decreases progressively as the epidemic advances, from July 16 to Nov. 16.

(x.) *Comparison of Mortality in the two Epidemics.*

The last, like the previous epidemic, extended over portions of two years; and the deaths, which were not very numerous in 1853, slightly exceeded the deaths in 1848. The deaths by cholera and diarrhœa in 1854 were, however, 5,000 less than the number that would have been told, had the epidemic been as fatal as the epidemic of 1849, allowing ten per cent. for increase in the population.

Years.	Deaths by Cholera.	Deaths by Diarrhœa.	Deaths by Cholera and Diarrhœa.
Year 1849 - - - - -	14,137	3,899	18,036
Deaths that <i>would have happened</i> if the population had been the same in the year 1849 as it was probably in 1854 - - - - }	15,587	4,299	19,886
Year 1854 - - - - -	10,806	4,000	14,806
Reduction of the mortality in 1854 as compared with that of 1849. - - - - }	4,781	299	5,080

The second outbreak began later in the season in the last epidemic than it did in the epidemic of 1849; and the

diminished figure of the mortality arises from the smaller number of deaths in the months of July and August. The disease had lost none of its virulence, and the deaths by cholera in September 1854, being 6,084, exceeded the deaths (5,031) in September 1849 by 1,053. The daily deaths by cholera in the two epidemics at the maximum, were 336 on September 4th, 1849, and 459 on September 2nd, 1854.

If it is considered that the interval between the epidemic of 1831-2 and 1848-9 was *seventeen* years, and that the epidemic which we have recorded followed the second epidemic after an interval of only *five* years, it is evident that the public apprehension for the sanitary state of London should suffer no abatement, but that the most active and complete measures should be adopted to prevent the approach or to mitigate the violence of impending visitations.

## SECOND SECTION.

### *Ætiology.*

#### A.—*Atmospheric Causes.*

IN reference to the *atmospheric conditions*, general or partial, which prevailed during the epidemic visitation, we have received reports, as follows, viz. :—from Mr. Glaisher, of the Royal Observatory, Greenwich, on the meteorology of London; from Dr. R. D. Thomson and Mr. Rainey, both of St. Thomas's Hospital, on certain chemical and microscopical investigations of air; and from officers of the Board of Health on the sanitary inspection of particular districts in the metropolis.

1. MR. GLAISHER'S REPORT is of peculiar interest. It presents the result of meteorological diaries kept, not only at ten stations previously existing, but at thirteen others specially established for the occasion; so that from the date of Mr. Glaisher's commission, observations more or less complete were made at no fewer than twenty-three sites in our vast metropolitan area. In reducing these multifarious observations, Mr. Glaisher's "first step was the examination of every reading in comparison with all others taken at or about the same time; the second was the application of index errors, corrections for diurnal range, and all necessary corrections and calculations to



deduce the mean daily value of each element of investigation. The weekly means of the daily values were next taken, and tables formed," exceeding fifty in number, as the basis of Mr. Glaisher's conclusions. From circumstances already adverted to, this admirable system of observation could not be completely organized till the epidemic had attained its climax; so that some gaps unavoidably remain in what, even with this deficiency, is a most valuable contribution to science.

Although Mr. Glaisher's report is in our Appendix, we think it needful here to present a short summary of its results; and we append a table which displays some of them in a compendious form:—

Deaths from Cholera.	Weeks ending as follows;	ATMOSPHERIC PRESSURE.			MEAN TEMPE- RATURE in excess.	HUMIDITY.		DENSITY of Atmo- sphere in excess.	No. of CALM DAYS.	Propor- tion of CLOUD, covered or (clear sky, 0).
		Total.	Aque- ous.	Total in excess, monthly mean.		Max. 100.	Excess, monthly mean.			
4	July 8	29°769	°368	-	-4°5	75	-	-	5	} 9
6	15	29°853	°388	-	-5°0	84	-	-	4	
33	22	30°114	°441	-	+1°3	72	-	-	3	} 5
180	29	30°151	°434	-	+4°1	68	-	-	3	
	JULY	-	-	+0°017	-	-	-5	+1		
488	Aug. 5	29°864	°429	-	-3°2	85	-	-	1	} 8
671	12	30°010	°402	-	-1°7	76	-	-	5	
772	19	29°968	°394	-	-1°3	74	-	-	6	
869	26	30°067	°398	-	+0°7	73	-	-	2	
	AUG.	-	-	+0°104	-	-	-39	+2		
1,646	Sept. 2	30°371	°455	-	+6°3	73	-	-	7	} 3
1,860	9	30°335	°371	-	+2°5	72	-	-	7	
1,527	16	30°014	°425	-	+4°1	80	-	-	2	} 6
1,182	23	30°115	°388	-	+2°7	80	-	-	0	
658	30	30°250	°342	-	+1°2	76	-	-	6	1½
	SEPT.	-	-	+0°199	-	-	-57	+2		
398	Oct. 7	29°878	°349	-	+2°1	80	-	-	3	6
227	14	30°198	°330	-	+1°0	83	-	-	2	4½
143	21	29°652	°280	-	-2°5	82	-	-	2	7½
48	28	30°685	°268	-	-2°2	83	-	-	5	5
	OCT.	-	-	+0°058	-	-	-16	+1		
25	Nov. 4	30°275	°319	-	+3°6	84	-	-	3	3½
16	11	30°275	°247	-	-1°7	83	-	-	1	6
10	18	29°625	°250	-	-1°0	88	-	-	1	8½
5	25	29°568	°215	-	-4°5	88	-	-	0	8
	NOV.	-	-	+0°003	-	-	+31	+5		
4	Dec. 2	29°686	°216	-	-2°5	83	-	-	0	6
2	9	29°810	°239	-	+1°5	83	-	-	0	5
0	16	30°029	°248	-	+3°1	84	-	-	0	6
2	23	29°767	°236	-	+1°8	87	-	-	0	7
0	30	30°125	°225	-	+1°6	86	-	-	0	5
	DEC.	-	-	-0°069	-	-	-17	-4		

Mr. Glaisher's inquiries have related to the *pressure of the atmosphere*, total and aqueous; to its *temperature*, mean and extreme; to its *moisture*, absolute and relative; to its *density*; to the directions and amount of its *movements*; to the *chemical and electrical* influences that act in it; to *haze, fog, mist, and rain-fall*.

(i.) The corrected weekly means of the observed *readings of the barometer* had been considerably in excess of their average during February, March, and April; but in the three months next following they presented no important deviation, and only became remarkable towards the end of August. The atmospheric pressure had then risen much above its normal amount, and during the worst period of the epidemic was more continuously great than at any other time. From the 25th of August to the 10th of September the reading was above 30 in.; and on three days in this period as high as  $30\frac{1}{2}$ . The mean reading for the two months exceeds the corresponding amount in any year of Mr. Glaisher's series; and it is the more noticeable since (as will presently appear) less than the usual effect was due to watery vapour.

(ii.) During the early part of 1854, the *mean daily temperature* of the air had been higher than normal; its excess, for the first 101 days of the year, averaging  $3^{\circ}\cdot4$ . There had then set in a very cold period, injuring vegetation and killing many hardy plants; and for the ninety-seven days terminating July 19th there had been a daily defect of temperature, averaging  $3^{\circ}\cdot3$ . The next few days showed a sudden increase of heat; the 25th of July was the hottest day of the year, its temperature rising nearly to  $90^{\circ}$ , and exceeding the normal by  $11^{\circ}$ . Three weeks of cooler weather followed; but from the 19th of August to the 11th of October (within which time were the worst ravages of disease) there was an excess of heat, averaging  $2^{\circ}\cdot6$  for each of the fifty-four days; and during one week of this period (that ending the 2nd of September) the excess amounted to  $6\frac{1}{4}^{\circ}$ . After the week ending the 14th of October, and excepting the week ending the 4th of November, the temperature was below its average till December.

(iii.) The *extremes of daily temperature*, and the range between them, have been noted by Mr. Glaisher carefully, and with some curious results. From the Greenwich observations it appears that, except June (which was

slightly in defect) every month of the year showed an excess above the average of diurnal range; March, April, and especially September being most remarkable in this respect; and the total result of this is, that, while for the thirteen years ending 1853 the mean yearly diurnal range was  $14^{\circ} \cdot 6$ , the range for 1854 was  $18^{\circ} \cdot 1$ , being  $3\frac{1}{2}^{\circ}$  above the average. But in comparing the extreme readings of the Greenwich and other outlying stations with those of London proper, Mr. Glaisher discovers the startling fact, that his central stations undergo a much less daily range of temperature; that, because of the dense veil which overhangs them, they, during day-time, cannot get equal heat from the sun, nor during night-time can equally cool themselves into space. Such excesses of night-temperature have amounted in the weekly mean to  $7^{\circ}$ ,  $8^{\circ}$ ,  $9^{\circ}$ , and  $10^{\circ}$ ; and as between particular stations, to  $15^{\circ}$  and  $20^{\circ}$ ; a period marked by this extreme difference having extended but for twelve nights' interval, from the 26th of August to the 4th of September.

Remembering that, amid the districts which most of all present this high night-temperature, there is spread the vast evaporating surface of the Thames,—a river which (so far at least as London is a drained city) represents the main sewer of our metropolis,—remembering, that from its putrescent banks and waters there arise vapour and miasm in proportion to that level of temperature, we must recognise the full right with which Mr. Glaisher insists upon this feature of our London climate. Almost uninterruptedly, too, the heat of the water is some degrees greater than that of the superincumbent air; for 28 consecutive nights, ending September 12th, this excess averaged  $16^{\circ} \cdot 3$ ; and there was another fortnight, beginning a few days later, during which it averaged  $16^{\circ} \cdot 5$ . At such periods we may (as Mr. Glaisher expresses it) infer the water to have been simmering, and the whole area of the Thames to have been giving off incessant and vast volumes of vapour, which, unsustained by the colder air, hovered over the city, thickened its atmosphere, occasioned the frequent prevalence of fog and mist, and explained the less daily range of temperature in stations overshadowed by its influence.

(iv.) The *vaporosity of the atmosphere* is estimated by a twofold standard; first, what quantity of vapour is actually holden in the air—how many grains per cubic foot—and what share does its weight contribute to the total of baro-



metric pressure? next, how near does that vapour approach to its own limit of maximum density—the limit at which (unless its temperature be raised) it admits of no further evaporation into it? An answer to the first question expresses an *absolute* quantity; an answer to the second, a *relative* quantity; and it is requisite to observe this distinction, because, on some damp wintry day for instance, the atmosphere may be within a shade of aqueous saturation, while in actual weight of water it scarcely holds the half of what, at higher temperatures, would leave it still capable of considerable drying-power. According to both standards, however, our atmosphere, during the chief prevalence of cholera, was less full than usual of aqueous vapour. In July, August, September, and October, it was further than usual from saturation; and from June to November it contained, in weight of vapour *per* given measure of air,  $\frac{1}{20}$ th less than its average.

(v.) In every month, excepting January and December, the *density of the atmosphere* has been in excess; the mean weight of a cubic foot of air having, for the year, been 2 grains above its average.

(vi.) With respect to the *movements of the atmosphere*, Mr. Glaisher's observations relate to the direction of winds, their mean force and velocity. From July 1st to September 11th the wind came alternately from S.W. and N.E. with nearly equal frequency, but with a difference of force greatly in favour of S.W.; in the next 28 days its direction varied more, but on 19 of them was W.S.W.; from October 11th it was W., and remained a compound of west to the end of the year. The daily motions of the air, irrespective of their direction, were much under the average; there was a defect in July of 34·3 per cent., in August of 25·7, in September of 15·3, in October of 29·3, in November of 32·8, from the respective averages of daily horizontal movement. Of the 123 fatal days, from July 1st to October 31st, there were 65 on which calm more or less prevailed; and it appears, as a fact of great interest in relation to the severity of the epidemic, that in the low-lying districts the air was at all times in much less motion than in those of higher level; entirely stagnant in the former, whenever in the latter it was noticed as calm; and when, at these, some hopeful wind blew with a pressure of 23 ounces, those suffering districts got but a sixth of the breeze.

(vii.) From July to the end of the year there were but



few *thunder storms* ; in fact, no great electrical disturbance took place from the time of the first outbreak of cholera in July, so long as the disease continued. Hail was noted on one day only within the same interval of time, viz. on October 23rd. So far, therefore, as the electrical observations indicate, in connexion with the much less than usual number of electrical disturbances in these months, it is inferred that there was a general deficiency in the tension of the common positive electricity prevalent during the period. No observations upon the electricity of the atmosphere were made till the disease was at its height ; at this time the electricity was positive but weak, and continued so till the end of September. Positive electricity, with tension somewhat greater than in September, was present at stations of moderate elevation, always except when rain was falling, in the months of October, November, and December. Common atmospheric positive electricity has therefore been as prevalent as usual. At stations situated nearly on a level with the river Thames, the electricity was generally weaker than at stations of higher elevation, and was more frequently negative.

(viii.) From August 24th till September 4th there was no *ozone* at any station near the metropolis, and very little at any station over the country ; a little was shown on September 5th, and from this time onward was general. At all stations of low elevation its amount has been insignificant, and at many near the river not a trace of it has been detected throughout the whole epidemic period ; while at places of high elevation it has nearly always been shown, and at intermediate stations occasionally ; seeming to graduate itself according to level, and to increase as we ascend from lower to higher ground.

(ix.) *Haze, fog, mist*, were singly or together prevalent on five days of July and eight days of August. The beginning of September was ushered in with a dense blue mist ; this in the second week of the month (when cholera was still at its height) was exchanged for a thick atmosphere of fog, which continued with little intermission till the end of the month, and at low places prevailed both day and night. During all this time, the distance was misty, middle-distance indistinct, and sunshine pale and watery ; but occasionally the atmosphere became partially translucent, and for awhile, in the higher levels of London, buildings would seem defined with remarkable clearness. The same kind of weather continued in October ; and

mist, fog, or haze, in or about London, was recorded on 19 days of November and 21 of December. The fortnight ending September 9th, and the week ending September 30th, were the periods, subsequent to July 1st, in which the sky was least overcast with *clouds*.

(x.) Of *rain*, there was a deficiency in every month of the year excepting May and December; and the whole year's amount was one quarter short of its average. Of 136 days following July 1st, there were 93 on which no drop of moisture fell, and 25 others with but very trifling rain. The crisis of the epidemic was in the driest period.

Mr. Glaisher has given additional value to his report by furnishing some particulars as to the meteorology of the epidemic periods of 1832 and 1849. In those years no observations were made at central points in the metropolis; and, therefore, it is only in respect of outlying stations that the meteorological phenomena admit of strict and detailed comparison. The records of 1832 relate to the atmospheric pressure and temperature, to the direction of the wind, fall of rain, clearness of sky, and frequency of electrical disturbances; those of 1849, except for their non-mention of ozone, admit of almost complete comparison with the present series. Mr. Glaisher's summary of these comparisons is in the following words:—

“The three epidemics were attended with a particular state of atmosphere, characterized by a prevalent mist, thin in high places, dense in low. During the height of the epidemic, in all cases, the reading of the barometer was remarkably high, the atmosphere thick; and in 1849 and 1854, the temperature above its average. A total absence of rain, and a stillness of air amounting almost to calm, accompanied the progress of the disease on each occasion. In places near the river, the night temperatures were high, with small diurnal range, with a dense torpid mist, and air charged with the many impurities arising from the exhalations of the Thames and adjoining marshes, a deficiency of electricity, and, as shown in 1854, a total absence of ozone, most probably destroyed by the decomposition of the organic matter with which the air in these situations is so strongly charged.

“In both 1849 and 1854, the first decline of the disease was marked by a decrease in the readings of the barometer, and in the temperature of air and water; the air, which previously had for a long time continued calm,

was succeeded by a strong S.W. wind, which soon dissipated the former stagnant and poisonous atmosphere. In both periods at the end of September, the temperature of the Thames fell below 60°, but in 1854 the barometer again increased, the air became again stagnant, and the decline of the disease was considerably checked. It continued, however, gradually to subside, although the months of November and December were nearly as misty as that of September. By the close of the year diarrhœa and cholera had subsided, but a high rate of mortality still continued.”

---

It now remains for us to appreciate, one by one, in their relation to life, the several meteorological deviations which Mr. Glaisher records. And this is no easy task; for the pathological meaning of many atmospheric variations, at least in their minor degrees, is hitherto quite unknown. What effect is produced on human life by an inch rise or fall in the barometer, by fluctuations of humidity and dew-point, even by seasons of non-average temperature, is very imperfectly measured. Still less is known of the chemical activities of the atmosphere. We know, indeed, that this boundless ocean of air is, in one-fifth of its volume, *oxygen*; the gas, which more or less rapidly brings all organic compounds into simpler chemical forms, exhausting those qualities that make putridity, and terminating those transitional states in which the powers of morbid infection resides. But there is every reason to believe that the oxidizing power of the air varies at different moments, as assuredly it varies at different spots. *Electrical discharges*—the frequent source of such variations—constantly occur in the atmosphere, developing in it, wherever it extends, that mysterious increase of its oxidizing power which is called *ozone*; a qualification of the air so subtle in its kind, that chemists still doubt whether it be a separate entity; yet in its function so definite, so hostile to organic miasms, so incompatible with them, that its presence enables us to affirm their absence; for wherever it meets them it must be spent in disinfecting their unwholesomeness, neither leaving of them any residue unneutralized, nor itself remaining free, except as predominant force have been on one side of this conflict or on the other. It is likewise probable that the great acts of aerial renovation are modified by the powers of *solar light*, wherein our world floats and revolves; since in every ray of it there are chemical



influenes, capable of affecting in turn each breath of the atmosphere they traverse. But these parts of the subject are of recent and unfinished discovery; our means of observation in regard of them are hitherto far from complete; and it would be premature to do more than point to such influenes, and to their possible fluctuations, as a field for future most important inquiry.

How requisite is such research, may easily be illustrated. Mr. Glaisher, in whose personal observations we should repose full confidence, informs us, that he can by sight estimate certain differences of mist, which he identifies with corresponding differences of epidemic sanitary condition; that he can connect one tint of mist with the prevalence of cholera, another with the prevalence of influenza; yet that, except for this rude test of colour, he cannot discriminate those mists, and has no hygrometrie or other meteorological knowledge of their existence.

Amid such uncertainties, we only venture to glance at the less obscure aspects of this interesting investigation.

The *undue height of the barometer* is an indication to which Mr. Glaisher draws particular attention, as having generally prevailed at the worst moments of each epidemic. During August and September 1854, it stood above its average from  $\frac{1}{10}$  to  $\frac{1}{2}$  in.; but lest undue importance should be attached to this one element, it must be noted, that in February (when there were but three cholera deaths) it had ranged nearly  $\frac{3}{10}$  in., and in March (when there was no such death) more than  $\frac{2}{5}$  in. above its average. We know of no direct influence which these atmospheric changes can have produced on human life, nor of any they can have exerted on the rate or kind of chemical change; but to a limited extent (as implying greater density of air) they would operate against vaporous diffusion, and in this degree may be probably estimated as favourable to the stagnation of miasm.

Of the immense *influence of heat* it is scarcely requisite to speak, in respect either of its well-known faculty to accelerate chemical changes, or of the many other differences that follow its range—the rarefaction of air, the lessening of humidity, the excitement of evaporation. Despite some exceptions, probably less real than apparent, it seems that Asiatic cholera, and indeed bowel-poisons generally, are favoured by high temperature; and in comparing together our two last epidemics, with a parallel comparison of their seasons, we are struck with the fact

that in 1854, when the summer temperature began later than in 1849, and quite abruptly rose to its maximum, so too the curve of mortality in that epidemic was peculiar,—seeming to imitate the summer temperature in its deferred commencement and sudden rise.

The *less range of London temperature* is a most important fact. It belongs not so much to our lesser heat by day as to our greater heat by night ; it means that London, though in the daytime somewhat less sunned than the outlying districts, sustains by night a considerable excess of temperature, with that more continuous activity of chemical decomposition which such an excess implies ; and this influence is the more important, as it is predominantly felt in those low alluvial districts, where the material for decomposition is most rife.

Of *fog, mist, and haze*, in their mere hygrometrical relations, we know nothing to affect life ; but it is requisite to remember that when these hang over districts of London—condensed in their ascent from the “simmering” river and filth-sodden soil—they represent not mere clouds of aqueous vapour, but, too probably, other products of terrestrial exhalation, delayed in their transit to space and withheld from the diffusion they had commenced.

The great *predominance of calm* was doubtless of baneful effect. It is a familiar matter, not only for households but for countries, that free ventilation is an indispensable condition of health. As individuals suffer their temporary inconvenience, when detained in unventilated rooms, and grow poisoned and cachectic if such be their habit of life ; so, in recesses of the earth’s surface, where, amid great mountain-chains, the ponded air lies unruffled by free whirls of wind, whole communities abort in the stagnant atmosphere, and beget a cretin, goitrous population. In respect of local impurities and the mischief they may engender, no miasm can survive an adequate commixture with air ; and in relation to the severity of epidemic disease, it is impossible to doubt that the more or less vehemence of aerial circulation is a variant of the utmost importance.

As leading results of Mr. Glaisher’s inquiry, two facts stand in relief:—

1, that the year 1854, and other years when cholera has prevailed, have had their marked meteorological characters, the general tendency of which has been to render

the season defective in those atmospheric changes which renew the purity of air ;

2, that these characters, apparently so definite in their meaning, are in their kind such as to prevail with greatly increased development in those low levels of London where all visitations of cholera have most cruelly pressed ; for high barometric pressure, excessive night-temperature and hazy air, with absence of wind, of ozone, and of electricity, would all (as the station-tables show) appear in their most marked degrees throughout those alluvial districts :

And, when these two statements are compared, it seems probable that in the atmospheric conditions of the year (or in some unknown influence essentially joined with them) there has been an important factor for the problem of that epidemic mortality.

In concluding our account of Mr. Glaisher's researches, we would submit one more extract from his paper:—  
 “ I have little hesitation in saying, that were the meteorology of our towns carefully ascertained and collated with that of the metropolis, and both together with that of the country generally (of which last I have a foundation of many years' continuous observations) that in a short time we should be in a condition to elaborate a clear insight into the meteorological causes of cholera, influenza, and many phases of disease which now burst upon us with the suddenness and devastating power of a divine and wrathful visitation.”

---

(2) SPECIAL EXAMINATIONS OF THE ATMOSPHERE were to some extent conducted, both chemically and microscopically, with the object of determining, in reference to the causation of the disease, whether any peculiar organic forms or unusual chemical products could be detected, either in the general air of an infected locality, or in its sewer-gases, or in the immediate atmosphere and exhalations of the sick.

The spot chosen for these observations was in the low-lying district of St. Olave, Southwark, where the epidemic had great prevalence, and where the wards of St. Thomas's Hospital gave the observers every facility for the requisite access to infected persons. The experiments were conducted by Dr. Thomson and Mr. Rainey, whose respective reports are contained in our Appendix, and whose method consisted in examining (the former chemically, the latter



microscopically) certain quantities of distilled water and of sulphuric acid, through which had been drawn, by a well-devised suction-apparatus, large volumes of the air which it was wished to investigate.—

From these examinations appeared as follows, viz.:—

(i.) In the atmosphere of a ward filled with cholera patients, while the disease was at its height, there were diffused various substances; some not possessed of life—the familiar dust of an inhabited room,—minute hairs, particles of smoke, epidermic scales, vegetable fibres of different kinds and colours, starch granules, &c.; others distinctly having life, and showing growth or movement. Of the latter, Mr. Rainey reports, that they had the appearance of small flocculent masses, visible to the naked eye, in the fluid in the bottom of the vessel; that, examined at the same time by Dr. Thomson and himself, they were found to consist of the mycelia of fungi, apparently in an active state of vegetation, mixed with the dusty impurities before mentioned; that he could discover no appreciable difference between these growths and the mycelia of fungi which had formed in solutions of vegetable substance after exposure to the air where no cholera was present; that besides the fungi, there were extremely minute, colourless, indistinctly beaded fibres (resembling in their general character that form of *Vibrionia* called *bacterium*) so abundant as to cover some of the larger branching fibres of the mycelium; and that these he does not recollect to have seen on mycelia growing in astringent vegetable solutions prepared for the purpose of producing fungi. Dr. Thomson adds, that this air gave very evident chemical signs of containing organic matter.

(ii.) In the atmosphere of a ward, only partially filled with cholera patients, when the disease was very much on the decline, the dead and living form, separated by filtration, were of the same kind as those just described; but the vibrionic fibres were much less numerous than in the former observation.

(iii.) In a third examination, made when cholera had left the district, and when, consequently, the ward was empty, dust particles were found, with the mycelia of fungi in considerable abundance, and apparently in active vegetation; but Mr. Rainey, with very careful search, could find no trace of the vibrionic forms.

(iv.) In the external atmosphere adjacent to the hospital, various dusty impurities were observed, as also sporules

and fungi to a considerable extent, but no vibriones. The collection of this air for examination began on the 21st October.

(v.) In air collected from within a sewer during twenty-seven days, beginning 22d November, there was found less dusty admixture than in the upper air. Vibriones were seen in much larger quantity than in any of the previous specimens, traversing the field of the microscope with great rapidity, and fungi were also present.

(vi.) The water through which sewer-gas had passed was strongly alkaline with ammonia, the result of organic decomposition; while the other specimens gave a powerful acid reaction, apparently from products of combustion contained in them.

The above results of a local examination have, at present, little more than a negative interest. The presence of fungi and their sporules in the atmosphere, appears to have had no relation whatever to the proximity of cholera patients, to have continued in the ward when vacated of all inmates, and to have been found in the exterior air (including that of the sewer) when cholera had long ceased to be epidemic in the district and in the metropolis. More importance might at first sight seem to belong to the presence of vibriones in the air, especially since they diminished when cholera patients were fewer in the ward, and vanished when no such patients remained. Yet, considering what is generally known of the habits of these peculiar forms, we cannot conclude that this was an essential coincidence. The development of vibriones is intimately related to animal decomposition; and the discovery of their profuse existence in sewer-gas, when cholera was no longer present, illustrates how they may multiply in an air that is loaded with organic miasm. These creatures are far too habitually about us for it to seem probable that they represent any new element in the causation of disease. We suspect that their diffusion in the cholera ward at the time of Mr. Rainey's first examination (when that ward contained its full complement of patients) simply indicated that the air was profusely charged with animal exhalation; that subsequently their existence varied with the number of persons occupying the ward; and that under similar conditions of temperature, ventilation, and cleanliness, they would have been equally abundant, though the inmates had been suffering from other disease.



We cannot pass this very interesting part of our subject without expressing our regret that researches of so much importance could not have been commenced at an earlier date, and been made more comprehensive in their scope. Many reasons will appear in the course of our report, to justify us in saying that a very complete and exact inquiry into the chemistry of organic decomposition during the epidemic prevalence of cholera—especially into the successive transformations of animal refuse at such times—might furnish all-important information as to the characteristic poison of the disease. It was with this belief that we suggested the necessity of submitting sewer miasm to examination; but from circumstances, quite beyond control, this part of the inquiry could not be undertaken till the time had passed for finding in it the solution we desired.

---

3. VALUABLE INFORMATION relative to local atmospheres has been gathered from other sources than direct chemical analysis. Nine sanitary inspectors were employed to visit all the localities most severely affected with cholera; and Dr. John Sutherland, late Superintending Inspector of the Board, has compiled from their statements an account, which is already before Parliament.\*

From Dr. Sutherland we learn that “all the inspectors agree in stating, as the result of their experience, that in those districts where cholera had become localised they found it connected with obvious removable causes”—these causes, so far as relates to the present subject, being such as affect the purity of the air, by loading it with the miasms of decomposing organic matter; that in considerable parts of the metropolis “there are large masses of population dependent for their drainage on open ditches, tidal ditches, old badly-constructed sewers, and still worse house-drains,—the result of the whole being that the *excreta* of a large part of the metropolis are not conveyed away, but are left to putrefy and rot in the open air, in cesspools under houses, or in large underground sewers, always generating foul gases, which are poured out into the streets or into the houses while in the more open districts the exhalations from the ditches keep the atmosphere in a con-

---

\* Letter of the President of the General Board of Health to the Right Honourable Viscount Palmerston, accompanying a Report from Dr. Sutherland on epidemic cholera in the metropolis in 1854; presented to both Houses of Parliament by command of Her Majesty.

stantly malarious condition;" that "the sewerage of all the localities is in a most imperfect condition;" that "a great sanitary evil incident to districts south of the river is the reflux of the tide through the sewers at high water, not unfrequently flooding the basement of houses, and in some parts rising in the gully-grate up to the level of the street;" that from the discharge of our metropolitan sewage into the river, "the banks at low water are in a filthy state" and "the exhalations from these pestilential banks of mud under a hot sun are most injurious to the purity of the air;" that "the pavement of courts and alleys is in general very defective, their gutters and those of the smaller and poorer streets badly constructed, retaining foul water before the doors of the houses;" that "the narrow back streets are very imperfectly cleansed, and the courts and alleys at ordinary times entirely neglected;" that bitter complaints are made of the non-removal of house refuse, which, with its various organic contents, is liable, for want of a properly organized system of dustmen, to remain as a nuisance for weeks; that in large districts of the metropolis there is "total absence of any comprehensive plan of laying out ground for building purposes, with a neglect of all arrangements for ensuring a free circulation of air round dwelling houses, and the existence of numerous narrow overcrowded courts and alleys, many of them mere *culs-de-sacs*;" "that in all the poorer classes of dwellings the means of ventilation are defective in the highest degree, the accommodation wretched and confined, the houses overcrowded and badly lighted, intermingled with private slaughter-houses, cowhouses, stables, pigsties, and public privies;" that "cellar dwellings are still in use in many parts of the metropolis;" that "the influence of noxious trades and nuisances in predisposing to attacks of cholera has been observed during the late epidemic;" dust-contractors, bone boilers, and gluemakers being specially mentioned, with "the effluvia proceeding from filthy stables, cowyards, and pigsties," and instances being referred to where whole families, exposed to such agencies, have been attacked, and in part fatally, by choleraic disease.

In contrast to the above characteristics of places in which cholera has prevailed, Dr. Sutherland refers to the condition of certain spots that have enjoyed a disproportionate immunity from the disease, apparently in result of sanitary amelioration. He cites instances where, at low levels in the metropolis and amidst a general infection of

the district, such an exemption has been enjoyed by particular groups of houses, which in former visitations had shared an equal lot with the rest, but which in the interval since those visitations, at the suggestion of a fatal experience, had had removed from within their atmosphere certain definite sources of contamination.

In the same sense it is reported that in common lodging-houses, formerly (by reason of overcrowding and filth) the fields of rich harvest for choleraic death, there has now, under that system of police-regulation which is due to Lord Shaftesbury's Act, appeared so little evidence of such infection, that Dr. Sutherland thinks it "necessary to make some allowance for imperfect information on the subject."

Similarly, among 2,791 persons who, at various points of the metropolis, were living under the auspices of societies for improvement of the dwellings of the poor, only six deaths by cholera occurred; of which one seemed due to an exterior infection, one to a poisonous supper of stale crab, and the remainder (which were all in a single family) to some cause evidently of most limited operation.

The cases of Bethlem Hospital and the City House of Occupations, on which Mr. Lawrence has favoured us with a memorandum (Appendix, No. V.) give their share of testimony to the same effect: these well-regulated establishments, with their population of 700 persons, having suffered no death from cholera while it prevailed most intensely around them.

The instance of the city of London, with 130,000 inhabitants, suffering in the late invasion 71 per cent. less cholera mortality than in 1849, is another illustration which Dr. Sutherland might have adduced of the epidemic lessening its pressure on a population in proportion as the sources of atmospheric impurity have yielded to sanitary improvement.

### B.—*The Water-supply of London.*

INQUIRIES into the water-supply of houses and districts suffering from cholera have been conducted under two heads, respectively *Chemical* and *Microscopical*.

1. The first of these, entrusted to Dr. Thomson, has given results, the sum of which, as regards our supply by the great trading companies, is embodied in the following table:—



GRAINS of ADMIXTURE per Gallon in WATERS of the under-mentioned COMPANIES.

	I.	II.	III.	IV.	V.	VI.	VII.	VIII.	IX.
	Lambeth Company.	Grand Junction Company.	West Middlesex Company.	Chelsea Company.	Southwark and Vauxhall Company.	New River Company at New River Head.	New River Company supplied to Soho, from a well near the Hampstead Road.	East London.	Kent Company.
1. Organic Matter	1·390	1·920	2·080	5·410	3·640	2·330	1·980	1·940	1·480
2. Silica	·350	·090	·520	1·511	·240	·180	·780	·320	·420
3. Sesquioxide of Iron, Alumina, and Phosphates	·215	·730	·460	·639	·460	·400	·210	·520	·130
4. Carbonate of Lime	10·144	8·870	9·919	9·550	10·700	11·985	7·085	11·997	9·540
5. Sulphate of Lime	2·149	2·368	2·109	6·432	3·179	1·326	1·523	·897	3·085
6. Chloride of Calcium	—	—	—	—	1·363	—	—	—	—
7. Nitrate of Lime	trace	trace	trace	trace	·076	trace	trace	trace	trace.
8. Carbonate of Magnesia	0·592	0·720	·720	·438	0·500	·855	1·185	·743	·210
9. Carbonate of Soda	—	—	—	—	—	trace	4·909	—	—
10. Sulphate of Magnesia	—	—	—	1·390	—	trace	trace	trace	trace.
11. Chloride of Magnesium	·617	·542	·360	1·947	2·101	trace	—	·237	·949
12. Sulphate of Potash	·730	·553	·577	2·903	2·413	·884	1·266	·682	1·153
13. Sulphate of Soda	—	—	—	—	—	—	8·051	—	—
14. Chloride of Sodium	·966	·947	1·637	29·797	16·001	2·355	7·807	1·125	·874
Total	17·153	16·740	18·443	60·017	40·673	20·315	34·796	18·461	17·841
Residue by Evaporation	17·440	16·920	18·970	60·170	41·780	20·780	35·050	18·300	17·760
15. Carbonate of Ammonia	·064	—	—	—	·840	—	—	—	—



In regard of the mineral impregnations shown in the above analyses, two features deserve particular notice; first, in all the specimens, that large quantity of lime-salts to which London water owes its costly and inconvenient property of *hardness*; secondly, in the two waters (IV. and V.) which are derived from the Thames at Battersea, the remarkable evidence given by chemical analysis, that an admixture from the sea reaches to this distance inland, and renders the river *brackish* with chloride of sodium.

Much significance belongs to what is set forth in the first line of the table—the proportions of *organic matter* dissolved in the several waters. Here great differences present themselves; for, as we compare together, in the first five columns, those waters which are drawn from the Thames, we observe that the admixture of such matter increases from 1·39, where the source of supply is at Thames Ditton, to 5·41, where it is at Chelsea. With the fact of this increase we remark that other materials, nitric acid and ammonia, have simultaneously risen into sensible quantity,—materials which are derived from the conversion of animal products. Without anticipating the demonstration presently to be founded on the microscopy of these waters, and even setting aside our knowledge of where the sewers disgorge into the river, it is already easy, on chemical evidence, to say that the Southwark and Vauxhall and the Chelsea Companies, pump their supply from a source profusely contaminated with the refuse of animal life.

With respect to water-supply generally, nothing can better prove a liability to foreign admixture than any *uncertainty of composition*; and Dr. Thomson, having for the most part made repeated analyses of waters severally purporting to be one in kind, observes, that the supply of each company, examined at different times, shows extensive fluctuations of impurity. The following table exhibits the range of such differences, as noted by him; and he adduces these facts to explain how widely his own report differs from that\* on which was founded the legislation of 1852,—the latter (he says) deducing its conclusions from samples of water drawn under too favourable circumstances, and strangely representing, as least contaminated with foreign

---

\* Report on the Chemical Quality of the Supply of Water to the Metropolis, by Professors Graham, Miller, and Hofmann; presented to Parliament by Her Majesty's Command. 1851.

matter, those which were taken nearest to London and most under influence of the tide.

GRAINS *per* GALLON of FOREIGN ADMIXTURE in the under-mentioned COMPANIES' WATERS.

Lambeth.	Southwark and Vauxhall.	Chelsea.	New River.	East London.	Kent.
12·12 to 17·98	22·50 to 72·66	36·96 to 65·66	15·75 to 35·05	17·02 to 19·60	15·02 to 21·10

2. The microscopical examinations, conducted by Dr. Hassall, form a necessary supplement to the chemical inquiry; and in referring to his results we will only premise that water, scrutinised with the highest magnifying power, reveals, if pure, no visible shapes whatever, and that consequently all such shapes discoverable by the microscope do, in their several kinds and grades, constitute a foreign impurity.

Having examined many specimens of water obtained from houses wherein one or more of the occupants had suffered from cholera; such water being supplied by the Southwark and Vauxhall, the Lambeth, the Kent, the East London, the New River, the Grand Junction, and the Chelsea Companies; Dr. Hassall reports among his results—

That the whole of the numerous specimens of water subjected to examination contained organic matter, dead and living, animal and vegetable; that the quantity and kinds of organic matter varied considerably in different cases, but were usually more or less constant for the same water; that the waters which contained the greatest number and variety of productions, dead and living, were from houses supplied by the Southwark and Vauxhall Company, that these abounded in living animal and vegetable forms of different genera and species; containing also a large quantity of dead organic matter, amongst which were frequently to be detected fragments of the husk of wheat, hairs of the same, starchy matters of different kinds, cells of potato and other vegetable tissues, with, in some cases, fragments of altered muscular fibre—these latter structures and elements being undoubtedly derived from the fæcal

matter contained in the sewage ; that the same species of organic productions were present in the waters obtained from houses supplied by the Chelsea Company, although in greatly diminished numbers—a result which might have been expected, since this company derives its supply from the same part of the Thames as the Southwark and Vauxhall Company, but filters it before delivery ; that the water procured from houses supplied by the New River, Grand Junction, and East London Water Companies, but especially the two former, all contained a great many organic productions ; that the water obtained from houses supplied by the Lambeth Company contained fewer organic productions than any of the rest ; that amongst the productions present in the water of the Southwark and Vauxhall, the Chelsea, and the West Middlesex Companies are several which are found only in brackish waters ; that the specimens taken from cisterns supplied by the Southwark and Vauxhall Company were very impure indeed ; far more so than any of the other waters examined, some of them being demonstrably contaminated with faecal matter.

Most of the above expressions describe the respective waters as drawn for domestic consumption, from the various butts and cisterns to which the companies supply them ; and as these receptacles for the most part favour the development of infusorial life, at the expense of dead organic matter, it may be assumed that the same waters drawn from mains or service-pipes might have presented larger quantities of dead tissue or excrement, and less maturity of living organisms. It hardly needs to be remarked, however, that butts and cisterns (however objectionable may be their use) cannot originate animal or vegetable growth ; and the derivation of such products is made obvious by Dr. Hassall's statement of what he observed in proceeding to the very sources of supply, and examining the waters of the Thames, the New River, and the Lea ; viz., that organic matter, both dead and living, animal and vegetable, was present in very considerable amount in the whole of these waters ; and that the living forms were discovered in considerable numbers, not merely in the deposit, but in nearly every drop of each of the waters, after they had stood the usual time, and after all but the lightest solid matter and the most active living infusoria had consequently subsided.

3. Both Dr. Thomson's and Dr. Hassall's inquiries have



extended to the examination of many well-waters in and about London.

In respect of superficial wells—those common sources of pump-water in the metropolis and elsewhere, the testimony now given strongly corroborates all that has frequently been urged as to the dangerous nature of such a supply. Both observers, though from different points of view, discover in these waters just such qualities as might be expected, from their having filtered through a porous soil, full of organic impurities,—that they contain sometimes evident sewage matter, sometimes an abundance of nitrates or of ammonia derived from the decomposition of animal substances, sometimes a variety of those animal and vegetable organisms which attest the progress of decay.

The deep well-waters, in respect of organic contamination, are usually in strong contrast to these; and in many of the specimens examined by Dr. Hassall there were no traces, or barely any, of infusorial life. Where such were found, their presence was to be accounted for by reference to special circumstances; by a communication of the well with some adjoining pond, or by its having had refuse wilfully thrown into it. But for these influences and the like, Dr. Hassall believes, “that scarcely a single organic production of any kind would have been found in any one of the (deep) well and spring waters subjected to microscopical examination.” The mere absence of such productions does not in itself establish the fitness of water for drinking, since there may remain various mineral admixtures to render it inconvenient or unwholesome; but both Dr. Thomson and Dr. Hassall refer with praise to the case of Woolwich, as one where a large population is supplied with water which presents an entire freedom from infusorial life, and which, though directly derived from the chalk formation, is artificially softened to about half the hardness of our Thames supply.

---

FROM such results, chemical and microscopical, as we have quoted from Dr. Thomson's and Dr. Hassall's reports, still more from their details, for which we must refer to documents printed in our Appendix (Nos. VII. and VIII.), it is evident that the commercial water-supply of London is derived from impure sources.



That furnished by the Lambeth Company is the best. Being taken from the river at Thames Ditton, it fulfils the requirements of the Metropolitan Water Act, and illustrates what after the present year will be supplied, with the sanction of the Legislature, to the greater part of the Metropolis. It is the best, but it is not good. Even in it Dr. Hassall finds "in not inconsiderable numbers, organic productions dead and living, animal and vegetable;" even in it Dr. Thompson finds traces of nitric acid, enough ammonia "to indicate an intermixture of sewage," and such proportions of organic matter as ought not to be insignificant to an educated community.

After the Lambeth Company follow, in the order of deterioration fixed by Dr. Thomson's analyses, the Kent, Grand Junction, East London, West Middlesex, New River, Southwark and Vauxhall, and Chelsea Companies. The last two greatly surpass the others in badness, and between themselves there is this difference:—while both draw from that part of the river, where the water is brackish from marine tides, and where an immense infusion of sewage proceeds uninterruptedly, the Chelsea Company seems to have the worse source for collecting, the Southwark and Vauxhall the lesser care for distributing its supply. Whether because of some greater influence of the tide along the northern side of the river at Battersea, or because of more sewage being discharged on that than on the southern side, the Chelsea water shows a much greater amount of dissolved impurities, but (apparently as the result of filtration) far fewer visible forms; while in the Southwark and Vauxhall water this evidence of unfiltered contamination reaches its highest degree, revealing to the microscope, not only swarms of infusorial life, but particles of undigested food referable to the discharges from human bowels.

It likewise appears from the evidence before us that the superficial wells of London afford a supply which, though often preferred for beverage in houses subject to the payment of water-rate, is generally not superior in quality to that distributed by the companies, and is liable to an aggravated form of the same contaminations.

SUCH having been the qualities of water consumed in the various parts of London during the late epidemic, it remains for us to examine how far the consumption of

these waters may have influenced the severity of cholera in London.

With respect to the *mineral ingredients* set forth in Dr. Thomson's analyses—although every grain of such admixture represents a deviation from the absolute purity of water—we have no reason for ascribing to them any appreciable influence in the matter adverted to. Considerable quantities of these mixed salts might be taken at a dose without producing serious injury to health; and we have no knowledge that, from their habitual consumption in small daily doses, there results any such chronic ill-effect as we might consider a probable predisposition to cholera.

With respect to the *living animal and vegetable forms* traced by Dr. Hassall through the whole series of waters, there seems no evidence that they, by their own action on the human body, could be productive of choleraic symptoms. There are indeed many instances, human and brute, of disease engendered in the living body through the tenantry of parasitic organisms, animal and vegetable; and, for aught we know to the contrary, many of the creatures described by Dr. Hassall may be capable of sustenance and multiplication within the bowels of those who swallow them. But in every known case where it can fairly be presumed that parasites are the causes of disease, they exist as a palpable morbid product occupying some considerable share of the affected body. The silkworm destroyed by *muscardine* dies because its whole body is riddled with parasitic vegetation, so dense that at last a mere heap of mould remains in place of the absorbed and disorganized animal; and every molecule of that mould makes evident the nature of the destructive process. What we know of parasitic diseases in the human subject—of hydatids and porrigo, for instance—tends all to the same point: in whatever way the foreign occupant have proved hurtful, whether it have starved the proper substance of the body on which it was grafted, or have provoked particular textures to acts of inflammation, or have choked their functions by its pressure,—itself, the causative thing, remains as a material shaped body, susceptible of ocular demonstration, side by side with its effects, and having bulk proportionate to them. Analogy would, therefore, lead us to infer that parasites could produce no attack of exhaustive purging and vomiting, except by having first

along the digestive canal multiplied to such swarms that they would be obvious to the most casual observer, both in the discharges of living patients and in the subsequent examination of the dead.

So far as this argument renders it improbable that the metropolitan water-supply produced cholera by means of the infusorial life engendered in it, the conclusion is strengthened by further remarks of Dr. Hassall's:—"Of the (living) organic products discovered in the waters examined, the great majority belonged to species which are known, and which have been long described in systematic works; and since the greater number of these are present in these waters at nearly all seasons, and since they are, therefore, constantly consumed, it is clear that they are in no way concerned in the production of cholera;"—"the number of those, the names of which were not known or their nature undetermined, was not considerable; and of these there was no one common to all the waters obtained from cholera houses which could be supposed to be influential in any degree in the production of the epidemic."

We do, however, attach very great importance to the fact, that nearly all the waters consumed in London, show a remarkable aptitude to develop low forms of animal and vegetable life; but this importance belongs, in our judgment, not to any direct influence exerted by such organisms on our own, but to the indications which their development affords that the waters wherein they grow are fraught with dead organic impurities.

The admixture of *decomposing organic matter* in the water-supply of the metropolis being attested equally by chemical analysis and by the microscopical evidence just adduced, we do not hesitate to speak of this contamination as one that may have exercised great influence on the spread of cholera among the population. The general history of this disease establishes its infinite preference for localities that are foetid with organic impurity; and it is impossible to conceive either any specific chemical changes arising in the air of a district, or any morbid action excitable by it in the living body—such changes or such action being due to its contamination by dead organic admixture—without recognizing that the water of the district likewise—great solvent of air as it is—must, if similarly polluted, be liable to undergo the same alteration, and to originate the same effects, as those of the atmosphere around it.



The present state of scientific knowledge does not justify dogmatic assertions on this subject ; but there are reasons for believing, in respect not only of cholera, but of many kindred diseases, that the means and agencies of morbid infection stand in intimate relation to decaying animal products within and without the body ; and the slightest taint of organic decomposition within the drinking water of a large population, therefore, constitutes a danger, which we cannot but regard with as much alarm as disgust.

---

If, in concluding our present section, we review the discussed materials in their bearing on the general ætiology of cholera, we think that the facts recorded enlarge the basis of previous knowledge.

The doctrine of epidemic cholera which has gained almost universal acceptance, does not affect to explain what may be that power—the exciting cause of the epidemic manifestation—which at intervals of time has forayed from place to place about our globe, sometimes vaguely spreading over a widened area, sometimes seeming to move in more defined procession, and which now for the third time has shed its fatal influence on our land. But with this mystery still unsolved, there has grown more and more into shape a doctrine which is both intelligible and practical ;—that the undiscovered power in its wanderings acts after the manner of a *ferment*, that it therefore takes effect only amid congenial circumstances, and that the stuff out of which it brews poison must be air or water abounding with organic impurity. Taking this as hypothesis, and testing it by the facts before us, we find that it would include and explain them.

Either in air or in water, it seems probable that the infection can grow. Often it is not easy to say which of these media may have been the chief scene of poisonous fermentation ; for the impurity of one commonly implies the impurity of both, and in considerable parts of the metropolis (where cholera has severely raged) there is rivalry of foulness between the two. But, on the whole evidence, it seems impossible to doubt that the influences, which determine in mass the geographical distribution of cholera in London, belong less to the water than to the air.

In our statistical section it appears that gradual exemption from the epidemic mortality “ has more nearly followed the degrees of elevation of soil than been proportionate to



any other general influence we could measure." But in this lessening scale—varying from a death-rate of 156 beside the river to one of 10 at the highest levels of London—there are so many and so considerable exceptions, as to show that the low level invites the epidemic invasion by reason of some attribute of its position which may elsewhere equally abound. Such an attribute is the excess of organic impurity, which (from the relation of the river to our London drainage) habitually saturates those alluvial parts; but which, through sanitary mismanagement or personal neglect, may prevail against the intentions of nature at the highest levels in the land.

Even among the best-placed parts of London there are districts so habitually foul and unwholesome, that a spectator might wonder whether by any result of tidal drainage the southern flats of the Metropolis can have been rendered more foetid and poisonous than they. But the meteorological history of London here comes to our aid; explaining how, even if there be equal filthiness in all districts, the poisonous consequences of filth must be inverse to the elevation of soil. For on the supposition (which this result greatly confirms) that the choleraic infection multiplies rather in air than in water, meteorology explains how the balance of healthfulness is weighted in favour of the higher levels, by their less participation in the high night-temperature of the metropolis, by their comparative clearness from mist, and above all by the curative resources of more free ventilation.

---

*Supplementary to the Second Section.*

REMARKS ON THE OUTBREAK OF CHOLERA IN SOHO.

ONE local outbreak of the epidemic—that which befel the neighbourhood of Berwick Street—was of such severity as to suggest that some especial causes must be concerned in its production. The circumstances of this remarkable attack have been investigated in a house-to-house visitation of the affected locality; and the results of that inquiry are stated in a Report (App. No. IV.) by the three gentlemen who conducted it—Dr. Fraser, Mr. Hughes, and Mr. Ludlow.

In the three registration sub-districts of St. Anne's, Golden Square, and Berwick Street, together comprising a population of 42,000 persons, it appears that there

occurred 537 deaths from cholera ; being at the rate of 128 to every 10,000 inhabitants, while the general cholera rate of the Metropolis was only sixty to the same number.

This high mortality was the more remarkable, as the affected districts are not situated at a low level, nor disproportionately inhabited by a poor population.

A striking feature of the outbreak was its extreme suddenness, as measured by the large number of persons almost simultaneously attacked. Its greatest local diffusion appears to have been reached on the second, if not on the first day, from its commencement ; it remained of equal prevalence for two days, and on each of the two following underwent a decline of 50 per cent.

In respect of this explosive appearance of the epidemic, it deserves particular mention, that for some time the district had been enjoying an exemption from disease quite out of keeping with its sanitary condition. Influences, universally recognised to be causes of disease, had been present, but in a manner for which we are unable to account had remained almost inoperative ; so that, till the very eve of this dreadful outbreak, the district might have boasted itself as one of average healthiness.

This fact is one of so much interest and importance that we have thought it advisable to have it thoroughly investigated by an inquiry into the mortuary statistics of the last seven years. The death-rate of the district during this period, if we could exclude from it those few days of epidemic visitation, would have been only  $20\frac{1}{2}$  per 1,000 per annum ; which, though far above what sanitary science can ensure to well-regulated districts, is considerably below our average metropolitan death-rate. Zymotic diseases, too, had made less than  $\frac{1}{2}\frac{1}{2}$  of this total.

It likewise deserves mention, that of the 537 cholera deaths of the late epidemic, 323 occurred in houses which, during the past seven years, had suffered no deaths from other zymotic disease ; and in comparison especially with the slight visitation of cholera which this district suffered in 1849, we discover that out of 86 houses in which cholera deaths then occurred, only 10 re-appear in the list of 310 houses in which cholera deaths occurred in the late epidemic. Since these results of our inquiry differ from what has been generally believed of the habits of the disease, we append (*see* p. 119) in detail the tables on which they are founded.

With respect to the causes of this particular outbreak, we find no apparent exception to the conclusions arrived at in the preceding section of our Report. Anticipating that for such an epidemic prevalence of cholera (whatever may have been the foreign influence that excited it) there must have pre-existed a certain local state of uncleanness with putrefiable matters, we trace no anomaly in the visitation.

That such local uncleanness prevailed most intensely throughout the suffering districts, is evident from the reported results of house-to-house visitation. The exterior atmosphere was offensive with effluvia from ill-conditioned sewers; the houses were almost universally affected in the same manner, partly from the same source, partly from their own extreme defects of drainage and cleanliness, partly from unregulated slaughtering and other offensive trades; the inhabitants were overcrowded, perhaps to the greatest degree known even in London; and the general architecture of the locality was such as to render it almost insusceptible of ventilation.

On the principles to which we have referred, and which we believe to be commonly recognised as presenting the most probable theory of choleraic irruptions, it will be obvious that the locality, notwithstanding its high level, contained every predisposing condition which (given the exciting cause) should render it prone to a violent epidemic explosion; and we believe that any person conversant with the laws of disease might have predicted its extreme liability to suffer what afterwards befel it.

Why, however, this district should have suffered in marked disproportion to many other districts, hardly, if at all, superior in their sanitary arrangements; or why, generally, it should be the tendency of cholera in its visitation to select particular foci for extreme outbreaks, instead of diffusing itself more equally over all ill-conditioned districts, is a difficulty which hitherto we have no scientific material to solve.

The meteorological conditions of the district were not minutely examined at the time of the attack: but the Reporters' mention of an atmospheric haze, and of a singularly stagnant, sultry, and oppressive air, leads us to believe that, if scientifically observed, they would have been found accordant with the generalizations in Mr. Glaisher's Metropolitan Report.



In explanation of the remarkable intensity of this outbreak within very definite limits, it has been suggested by Dr. Snow, that the real cause of whatever was peculiar in the case lay in the general use of one particular well, situate at Broad Street in the middle of the district, and having (it was imagined) its waters contaminated with the rice-water evacuations of cholera patients.

After careful inquiry, we see no reason to adopt this belief. We do not find it established that the water\* was contaminated in the manner alleged; nor is there before us any sufficient evidence to show, whether inhabitants of the district, drinking from that well, suffered in proportion more than other inhabitants of the district who drank from other sources.

There is mentioned, however, a remarkable instance in which it seems probable that the water of this well did really act as a vehicle of choleraic infection; but (assuming the absence of fallacy in the case) this probability might easily be admitted, without its thence resulting that infection depended on the specific material alleged. The water was undeniably impure with organic contamination; and we have already argued that, if, at the times of epidemic invasion there be operating in the air some influence which converts putrefiable impurities into a specific poison, the water of the locality, in proportion as it contains such impurities, would probably be liable to similar poisonous conversion. Thus, if the Broad Street pump did actually become a source of disease to persons dwelling at a distance, we believe that this may have depended on other organic impurities than those exclusively referred to, and may have arisen, not in its containing choleraic excrements, but simply in the fact of its impure waters having participated in the atmospheric infection of the district.

---

### THIRD SECTION.

---

#### *Practical Pathology.*

ONE of the earliest duties which devolved on us, as a committee for the scientific purposes of the Medical Council, was to suggest means for collecting and utilising

---

\* Dr. Thomson and Dr. Hassall examined this water, though not at the moment of the outbreak, and their account of it will be found in their respective reports.



the results of detailed medical experience in its individual circles of practice. For it was hoped that, in comparing together the contributions of many independent observers we might educe trustworthy information on various disputed particulars.

We accordingly prepared for circulation among members of our profession the Forms (A. and B.) already adverted to. These were so constructed that the returns, when complete, would inform us not only as to the stages of the disease, their duration, fatality, and relative frequency, but likewise as to each respondent's personal experience in the application of different remedies.

We now account in full for the mass of valuable information received in reply to this appeal. All that related to the history of the epidemic, irrespectively of medical treatment, has been extensively used in the statistical section of our Report, and has furnished material for our first series of Tables. The remainder, recording the therapeutical experience of more than 300 contributors, and representing in this respect an unprecedented multitude of authentic evidence, was deemed of sufficient value to justify its being referred to a special section of the Medical Council, whose report on the results deduced from it is already before Parliament and the public.

There was another class of contribution which we were most anxious to encourage. However painful the truth, we could not disguise from ourselves, as to the advanced stages of cholera, that nearly all resources of medicine seemed equally powerless for good; that practitioners, summoned to the relief of this dreadful disease, could scarcely decide between any two lines of treatment except for an occasional certainty that one of them would positively harm; and we felt deeply assured that—failing some happy chance of empiricism, there could be but one escape from this helpless condition of our art. We believed that the varieties and irresolution of practice corresponded to the actual unripeness of pathological knowledge, to the absence of those scientific principles which alone could give secure and uniform guidance: for, how futile to discuss the hopefulness of this or that experiment in treatment, while our profession was still unfixed as to what action in the suffering body it should be the definite aim of medicine to effect! If better success was to be won, we thought it must be through stricter studies in the nature of the disease, through learning as it were the mechanism of

its production, and appreciating, fact by fact, the full significance of all its symptoms.

During the preceding epidemic, both here and especially on the Continent, researches of great interest and importance had been carried on. Yet a vast deal remained for inquiry ; and we hoped it might be conducive to good, that this state of the case should be represented to persons who were likely to take part in scientific investigations.

We accordingly suggested "that there still remain in the pathology of the disease many undetermined particulars ; and that in respect of these, great assistance would be given if the physicians of London hospitals, and other officers of public institutions within whose province it falls to be cognizant of such matters, would direct their attention to specific pathological questions, and would communicate to us the results of their inquiry." Being then requested to specify those problems in the disease which we considered most deserving of study, we prepared a *Memorandum* on the subject which was transmitted to the chief medical officer of the various hospitals and infirmaries in the metropolis.

The epidemic, however, had already passed its climax, and soon rapidly declined and ceased. There was consequently little time or opportunity for new scientific observations ; and although some of the problems proposed might have been elucidated by the results of experience already gained, no replies were elicited from public institutions, except *St. Bartholomew's*, *St. Thomas's*, and the *Middlesex Hospitals*, and the *City of London and East London Dispensary*. Nor, independently of our appeal, did the practical pathology of cholera during its recent visitation appear to attract any active scientific research. In the epidemic of 1848-9, new methods of investigation had been most zealously employed ; and in 1853 the subject had lost its novelty, even for the younger cultivators of pathological science.

Reports bearing on different points of the inquiry were, however, received from Dr. R. D. Thomson, Dr. Hassall, and Mr. Rainey. These are printed in our Appendix, either entire or in part ; and their contents, as well as other facts communicated to us, will be noticed here only so far as they tend to elucidate those questions to which our *Memorandum* had called attention.

1. The solution of the first question—*Through what channel does the exterior cause or poison of cholera first*

*enter or affect the human body? is it through the lungs? or through the stomach and intestines? or otherwise?* obviously might be aided by evidence of various kinds; for example—1, by facts showing whether the atmosphere of localities infected by cholera has, or has not, properties favourable to the existence and increase in it of an organic or inorganic poisonous matter; or whether it contains or does not contain some matter of a peculiar character which might be regarded as the poison itself; 2, by evidence of the presence of such peculiar matter in the air-tubes of the lungs; 3, by facts demonstrating, on the other hand, its presence in the water drunk or in other articles of diet; and, 4, by the demonstration of its existence in the stomach and intestines, or in the discharges from them.

Some of the reports laid before us contain information relative to these several points.

(i.) The important observations of Mr. Glaisher, already noticed at length, demonstrate that not only in 1854, but likewise in 1849 and, as far as can now be ascertained, 1832, many conditions of the London atmosphere were such as would favor the retention and increase of any poisonous matter in the air, and that these conditions increased concurrently with the rise of the epidemic, and diminished with its decline. Such facts accord well with the view that the poison of cholera enters the human body through the lungs.

Researches already mentioned, of Dr. Thomson and Mr. Rainey, on the chemical and microscopical qualities of the air, in infected districts and in the immediate vicinity of the sick, have failed to advance knowledge in this particular. As was stated in our last section, they discovered in such atmospheres as they examined no ingredient which, with any plausibility, could be considered special to the prevailing disease.

(ii.) In further search for such possible foreign elements, should they have entered the body by respiration, Mr. Rainey and Dr. Hassall examined microscopically the lining membrane of the air-tubes in persons dead of cholera. Mr. Rainey, having explored the windpipes of eleven persons, as soon as possible after death, "could detect nothing like spores or the mycelia of fungi, or the vibrio-like fibres in any one of them." Dr. Hassall found vibriones abounding in the mucus of the bronchial tubes, but remarks that their presence was probably due to incipient putrefaction.



The general result of these observations, then, is negative in regard to the presence of any peculiar bodies in the air or in the lungs. For, although vibriones were found very abundant in the water through which the air of an hospital ward was passed when it was full of cholera patients, and very few or none when the ward was empty, we have already shown that in these coincidences there is no proof of vibriones having any direct relation with the cause of cholera.

(iii.) That the exterior cause or poison of the disease may enter the human body through the stomach and intestines, has been conceded by many persons who believe they can refer the causation of certain cases to the consumption of foul water, or other putrescent articles of diet; and in our last section we stated the reasons which would lead us to regard such consumption as a probable risk of empoisonment. But, as regards the most possible means of such infection—the water-supply of the metropolis—neither Dr. Thomson by chemical inquiry, nor Dr. Hassall with the microscope, has been able to identify in it any matter, special to the epidemic time or to the infected localities. Indeed (as we have already quoted) the latter observer expressly states that the very various animal and vegetable living products which abounded in the waters of cholera districts belong to species which are well known, and are to be found in the waters of districts not visited by the epidemic; that they have no especial relation with the presence of cholera; and are important only as affording evidence of the impurity of the water.

(iv.) Again, in the intestinal discharges Dr. Hassall has discovered no sporules or threads of any species of fungus, and no peculiar body of any kind, other than vibriones. Vibriones, he states, are constantly present in extraordinary numbers, not only in the matters discharged from the body, but also in the fluids taken after death from the intestines themselves, as high as the duodenum. But these bodies exist in other morbid intestinal discharges, and according to Mr. Rainey, they may be found also in the contents of the intestinal canal after death from other diseases, even when the examination is made at an early period. Their extraordinary number in the rice-water discharges, therefore, probably shows merely a great proneness to decomposition in the fluids poured into the intestines in cholera. It is questionable whether there is any connexion between the presence of vibriones in such vast number



in the intestinal discharges and the fact observed by Dr. Thomson and Mr. Rainey that the same bodies are found in the water through which the air of a cholera ward has been passed. Dr. Hassall states that neither vibriones nor the sporules of fungi rise into the atmosphere through the mere evaporation of the fluid containing them. Yet, as in a cholera ward, portions of the intestinal discharges would certainly be spilled on the floor or on the bed linen and become dry there, vibriones contained in them might in this dried state be disseminated through the air of the ward. We have already stated our suspicion that, when the scope of such examinations is extended, the presence or absence of vibriones in the air, or in the water through which it is filtered, will be found to depend on other circumstances—on the degree of crowding in the ward, and on conditions of ventilation and temperature, which would affect the quantity of organic matter diffused, and the rate of its putrefaction. But all doubt on the matter might be soon solved by parallel examinations of air obtained from rooms equally full of non-infected persons.

In reference to the possibility of infection through the gastro-intestinal membrane, we may here mention an exclusive form in which the doctrine has been urged,—to the effect that this is the only channel through which infection can occur, and that its invariable means consist in the swallowing of matters (chiefly water) specifically contaminated by the fæces of previous choleraic patients. We cannot doubt that drinking-water fouled with excrements, whether diarrhœal or healthy, would represent in a high degree those qualities of organic taint, against which we have already in general terms expressed our opinion. But, while quite prepared to admit the danger of that *class of ingesta* to which such water would belong, we can find nothing in support of the exclusive theory adverted to; and we believe, as already stated, that the geographical distribution of cholera in the metropolis, at each of its three visitations, has in its main features expressed, beyond the possibility of reasonable doubt, that its diffusion chiefly depends on other than dietetic influences.

2. In reply to the second question, “*Has the disease a period of incubation? if so, how long? and on what is it contingent?*” Dr. M'Loughlin remarks, that, during the prevalence of the epidemic, all persons who do not actually labour under cholera or diarrhœa suffer from a

slighter disturbance of the stomach and bowels, or from an unusual susceptibility to the action of purgative medicine. Further information is desirable respecting the latter condition, which seems sometimes to be well marked. But the knowledge which the question was chiefly intended to elicit was, whether an interval of definite or indefinite length elapses between exposure to the exterior cause of the disease and the first manifestation of its symptoms.

3. With reference to "*the communication of the disease from person to person*," no large body of evidence, either of a positive or of a negative tendency, has been received. In the paper communicated from the Middlesex Hospital, it is stated that in that institution "no disproportionate liability to the disease has been manifested in those who have been engaged in attendance on the sick, or engaged about their dead bodies, or occupied in cleaning their linen." "One patient and two nurses were attacked while in the hospital." In St. Thomas's Hospital the staff of "sisters" and "nurses" consists of 19 sisters and 47 nurses. One sister, who had at first the charge of the special wards\* for cholera, and four nurses not in attendance in the cholera wards, passed through the different stages into complete collapse. The sister recovered; three of the nurses died.

Mr. Whitfield, the resident medical officer of St. Thomas's Hospital, expresses his disbelief in the communication of the disease from person to person, but says he has seen many remarkable cases of the occurrence of the disease after communication with persons affected at the time or recently. He instances two cases. In one he attributes the attack to the alarm excited in the patient by her being told, when suffering from bilious diarrhœa, that the person in the next bed had had cholera—consecutive fever at the time of admission: the bilious diarrhœa almost immediately became choleraic. The other was the case of a nurse who became very sick and faint while engaged in rinsing out the sheets and linen of a cholera patient, and was attacked with cholera which rapidly proved fatal. The sheets and linen had been left unwashed from the previous day, and Mr. Whitfield believes that this case is an illustration of Professor Liebig's remarks, and that the nurse "imbibed

---

\* It may be remarked that these wards are the least favourably placed parts of St. Thomas' hospital, and that the hospital itself stands in the middle of a district where cholera has always prevailed with peculiar severity.

“ the poison, not by communication with the person, but from the decomposed secretion.”

We have already referred to a speculation, so far as it relates to the human subject, that all diffusion of cholera depends on a deglutition of some amount of choleraic fæces, conveyed from the sick to the healthy in various articles of food, and especially in water. Professor Liebig had stated as the result of experiments, that the rice-water discharges acquired at a certain stage of putrefaction the property of inducing a disease similar to cholera, in animals to which they were administered. Dr. Thomson communicates the particulars of an experiment of this kind, instituted by himself. The result was negative. But Dr. Thomson does not regard this single observation as conclusive. Moreover, it cannot be expected that the question will be settled in this way. There is no sufficient ground for thinking that the lower animals are susceptible of epidemic cholera; and the fact of diarrhœa being produced in them by such experiments as those of Liebig, would merely prove that the discharges, when putrescent, are noxious, not that they contain the particular poison by which cholera is generated in the human subject.

Some microscopic observations of Dr. Hassall on the clothes of cholera patients have an indirect bearing on the question of the mode of diffusion of cholera. For, even though the cause of cholera be not an emanation from the bodies of the sick, if it be a matter which increases in foul air, it may possibly be conveyed in the clothes of men from one locality to another. Dr. Hassall found, as was to be expected, that animal matters in which living organic products were visible could sometimes be extracted from the clothes of cholera patients; but he found no peculiar organic body. Sporules of fungi were very rare, and vibrios were abundant only in clothes which were stained with rice-water discharges.

4. The seventh question suggested was, “ *Does cholera begin as a morbid process of the gastro-intestinal mucous membrane, or is this preceded by some state of general poisoning which requires the gastro-intestinal membrane to act as an emunctory? Is the state of collapse determined by this gastro-intestinal flux, and in proportion to it, or can it arise independently of any such flux? How are the lividity and the cramps determined and proportioned?* ”



To the first two clauses of this question no direct answer has been received by us; yet they are of very high importance. The flux from the gastro-intestinal mucous membrane is, doubtless, the first of the more obvious phenomena of the disease, and may perhaps be in every sense its beginning; but, on the other hand, it may be that the cause of cholera affects primarily some other part of the economy, as the blood or the nervous system, and acts only through them upon the mucous membrane of the stomach and intestines; and it may be that the affection of this membrane is essentially the process by which some poison is eliminated from the body. Both views have been asserted by writers on the disease; and as they almost necessarily lead to opposite methods of treatment, it was thought desirable by this question to invite closer examination of the point.

With respect to the next clause of the seventh question, —“ *Is the state of collapse determined by the gastro-intestinal flux, and in proportion to it, or can it arise independently of such flux?*” Some relevant information is found in the returns of cases of diarrhœa and cholera treated during the late epidemic. The number of cases of collapse reported is 1,798, and in all but six cases the collapse was preceded by one or more of the earlier stages of the disease, which are attended by intestinal discharges. More accurate inquiry would most probably have elicited the fact, that even in the six exceptional cases some intestinal flux occurred before the phenomena of collapse were developed. In some or all of the six cases the amount of liquid matter discharged may have been inadequate to account for the collapse, but, even then, there may have been poured out from the blood a large quantity of fluid which still remained in the intestines. The following is an extract from the report received from the Middlesex Hospital:—“ A child about  $3\frac{1}{2}$  years of age was admitted in a state of complete collapse, having had only one motion and having vomited but slightly since the commencement of the disease. In this, as in many similar instances, it may be asserted, that the flux had already taken place into the bowel, but had not been discharged *per anum*, for soon after admission the child was profusely purged.”

In the Middlesex Hospital report, it is stated that although in general the severity of the collapse was in proportion to the amount of discharges, it was not so in



all cases. This is in accordance with previous experience, but it needs to be determined by further observation, whether the differences in amount of the discharges producing a given effect, and the differences in effect resulting from discharges of given amount, can be referred to diversities in age, in nervous power, in tendency to syncope, in the quantity of the fluid in the body, or in other like conditions of the individual patients.

5. The eighth group of questions related to the pathology of the *consecutive fever*; and the two questions "*To what extent does it depend on the previous occurrence of profuse discharges? or on the completeness of collapse? Does stupor in this stage always depend on uræmia? or on what?*" are in some measure elucidated, or at least touched upon in the papers before us. The tabular returns of cases of diarrhœa and cholera having been drawn up for the most part after the cessation of the epidemic, of course could not be referred to with confidence for the settlement of questions relative to minute points of pathology. But there is no reason to doubt the trustworthiness of the information of a general character which they afford in regard to the connexion between consecutive fever and collapse. 874 cases of consecutive fever are reported, 249 fatal, and 625 recovered from. In all the cases the fever stage was preceded by cholera; but the presence or absence of collapse is mentioned in only 808 cases; and in 223 cases of this number, or 27·6 per cent., the cholera had not passed into collapse. Here, then, it appears that the occurrence of consecutive fever is not closely dependent on preceding collapse. But the result is very different when the 225 fatal cases, included among the 808 cases above mentioned, are examined separately: for, of these, 199 were preceded by collapse, and only 26, or not quite 11·15 per cent., by cholera without collapse; so that the severer forms of the morbid states included under the term consecutive fever, seem to be in a very large majority of cases connected with a preceding condition of collapse.

These deductions from the tabular returns are entirely consistent with the results obtained at the Middlesex Hospital, as they are stated in the subjoined conclusions extracted from the report from that institution:

(a.) "Nearly all those cases in which collapse was complete and purging profuse, fell into a greater or less degree of consecutive fever."

(b.) “Many cases passed into the most extreme stage of collapse, and yet, after remaining in that condition for many hours, recovered with the slightest degree of consecutive fever.”

(c.) “Of those cases which did not pass into a state of complete collapse, several were followed by a slight and two by a severe form of consecutive fever.”

(d.) “All the cases of severe fever, with the two exceptions above mentioned, were preceded by severe collapse and considerable discharges.”

(e.) “The degree of fever appears to have borne some relation to the duration of the stage of collapse; there are, however, numerous exceptions to this rule.”

With reference to the question—“*Does stupor in this stage (consecutive fever) always depend on uræmia? or on what?*” the following remarks occur in the Middlesex Hospital report :

“Of the cases which died in a state of coma, with one exception, all had suppression of urine; and in the exceptional case the quantity of urine was diminished.”

“Case 39, fell into a state of complete coma from which he appeared sinking. After remaining in this condition for more than 24 hours he recovered. Throughout all this stage urine was passed copiously, and of good specific gravity. And although the urine contained a trace of albumen (as was the case with most cholera patients), there did not appear to be any reason for considering that uræmia was present. The coma much more resembled that seen in typhus.”

6. The import, tendency, and characters of the diarrhœa, which prevails epidemically together with cholera, formed the subject of the ninth group of questions.

(i.) “*When diarrhœa and cholera prevail together epidemically are they (with differences of degree) the same disease?*”—This question must now, doubtless, be answered affirmatively. The larger part at least of the diarrhœa which is generally so rife in localities where cholera exists, must be ascribed to the same cause and must be regarded as only a slight degree of the diseased action which in a higher degree becomes cholera. But if diarrhœa when aggravated is cholera, how comes it that so many deaths are during a cholera epidemic registered as caused by diarrhœa? During the 18 months from July 1853, to December 1854, 11,661 deaths from cholera were registered in London, and as many as 6,258 deaths from

diarrhœa. Making an ample deduction from this number on the score of the deaths from bowel complaints of various forms which occur annually when cholera does not prevail, there remain 4,000 deaths referable to the epidemic and registered as due to diarrhœa. How did these 4,000 cases differ from the 11,661 cholera deaths? That many of them presented distinctive features cannot be doubted. In the first place the average duration of the 4,150 cases of fatal diarrhœa registered was 13 days; so that a large portion of them must have been cases in which the disease ran a slow course; and, in the second place, children and old persons above the age of 65 furnished a large majority of them. Now there are fatal cases of the disease in which the lividity of the surface and the cramps are either very little marked or are absent, and these cases are observed chiefly in children and old persons, and are often of a lingering character. They, therefore, in all probability, form a part of the cases which appear in the register of deaths as fatal cases of diarrhœa. In the hospitals of London few deaths were attributed to diarrhœa, only 42, while the deaths attributed to cholera were 800. In workhouses, on the contrary, 400 deaths from diarrhœa were registered, with 924 deaths from cholera. This difference is in part explicable by the small proportion of children and old persons among the patients admitted into the general hospitals, and the large proportions of persons of those periods of life among the inmates of workhouses. Again in the districts on the higher levels the registered mortality from diarrhœa was in proportion to the registered mortality from cholera far greater than it was in the districts on the lower levels. And this may in part be explained on the assumption that in the higher districts where the epidemic cause was not generally in strong action, the fatal cases would often be less intense in character and less rapid in their course. These cannot, however, be accepted as the sole reasons of the predominance of diarrhœa amongst the deaths from the epidemic in the higher districts of London and of the small proportion it forms amongst the deaths in the hospitals. For in some of the hospitals where the deaths from cholera were numerous, no deaths from diarrhœa were registered, and in some of the sub-districts of London which have not a high level and which suffered severely from the epidemic, the deaths from diarrhœa exceeded in number the deaths from cholera. The truth probably is that the choice of



the term used in registering the deaths was often made somewhat arbitrarily ; that in private practice, especially in districts where the disease was not very rife, there was a disposition to give the less formidable name of “diarrhœa” even to cases which had the features of cholera distinctly marked ; whilst in hospitals, a more strictly pathological view of the matter being taken, all, or nearly all, the fatal cases were denominated deaths from cholera.

The deaths registered, as caused by diarrhœa, so far as they were due to the epidemic cause, were, then, cases of cholera more or less modified in their features, and had no closer relation with the ordinary cases of diarrhœa which were not fatal than the deaths registered as “cholera deaths” had ; for both classes of deaths were alike the results of the same morbid process which in its slightest degree was the “simple diarrhœa.”

(ii.) On the mutual relation of these two grades of the epidemic influence, we have received a communication from Dr. M'Loughlin, who is known to have devoted much time and labour to the inquiry. He strives chiefly to prove that cholera is *always* preceded for a certain time, some hours, some days, or some weeks, by a “premonitory diarrhœa ;” and he does not admit the occurrence of exceptional cases. It may be doubted whether the single profuse discharge which in a few cases precedes for an hour or little more the state of developed cholera, can be regarded as constituting *premonitory* diarrhœa. But it cannot be disputed that these cases are very rare, and that cholera is in the vast majority of cases preceded by a stage of diarrhœa which affords time for treatment.

(iii.) No evidence has been laid before us which might aid to solve the question, “*Does the diarrhœa, if left to itself, generally and safely tend to spontaneous recovery ; or do such cases, without medical treatment, frequently in proportion to their numbers, pass into true cholera ?*” And it will always be difficult to obtain satisfactory evidence with regard to this question.

For although the number of cases of cholera which were neglected in the stage of diarrhœa may be learned with accuracy, it will be scarcely possible to ascertain the number of cases of diarrhœa in the same town or district which terminate by recovery in that stage although not brought under medical treatment. The small proportion of the cases of diarrhœa which pass into cholera when remedial means are employed is shown in the “Report on



the Results of the different Methods of Treatment pursued in Epidemic Cholera."

7. No new *chemical analysis of the blood* in cholera has been communicated to the Medical Council. But as the changes which the composition of the blood undergoes necessarily bear a close relation to the composition of the fluids poured out from the blood-vessels into the intestinal canal, reference may here be made to the Report of Dr. R. D. Thomson (App. No. X.) on "the Chemistry of the Rice-water Excretions."

Some observations on the physical and microscopical characters of the blood after death from cholera will be found in Mr. Callender's minute account (App. No. XIII.) of twelve autopsies made at St. Bartholomew's Hospital. Remarks by Dr. Hassall on the same subject, and on the state of the urine in cholera, are likewise contained in our Appendix (No. XI.)

In any future visitation of cholera, more extensive and systematic inquiries, it is to be hoped, will be instituted relative to the pathology of the disease; and in order that these inquiries should produce large results, it is most desirable that they should be commenced at an early period of the epidemic. On this account (*see* p. 127) we append to our Report the Pathological Memorandum, which elicited the various answers discussed in our present section; and which may at least serve to remind pathological inquirers at a future period of the position at which our knowledge had arrived at the time of the last epidemic, and of the directions in which its advance was then felt to be most needed.

---

In closing our present Report, and therewith terminating our labours as a committee for scientific inquiries in relation to the late epidemic, we would venture respectfully to suggest, with a view to future emergencies of the kind that if such inquiries are to be fruitful of result, they should to some extent be continued in the absence of the disease which they aim at elucidating.

Just as there can exist no science of morbid anatomy, till the structure of the healthy body be first well learnt; just as there can be no medical knowledge of what makes a symptom of disease, till physiology have first established a standard of natural function; so it is, we believe, with

the studies which, in their future development, may perhaps explain the birth of epidemic disease.

The preparatory steps of such studies are even yet scarcely made. In attempts to elucidate the exact nature of those processes by which epidemic poisons are generated, almost insuperable difficulties have been found; not from want of apprehension as to what the factors may be in such exceptional generation of poison, but from lack of definite knowledge as to the normal working of these factors.

It seems certain that in the chemistry of organic decomposition there is concealed a large share of the mystery we would solve; and it is impossible yet to say, how much of the remainder may belong to undeveloped branches of meteorological science.

Under any future epidemic invasion which our country may unhappily suffer, it is in these departments of natural science that temporary and exceptional phenomena will claim renewed investigation; and we hope we do not outstep a proper fulfilment of the trust reposed in us, when we suggest, that in this intervening time are comprised golden opportunities for rendering probable the success of investigations then to be undertaken,—opportunities for making effective progress in the preliminary studies to which we advert, and for establishing a better normal standard than is yet discovered, to measure the chemical and meteorological anomalies of an epidemic period.

We have the honor to be, Sir,

Your obedient humble Servants,

N. ARNOTT.

WILLIAM BALY.

WILLIAM FARR.

RICH<sup>d</sup> OWEN.

JOHN SIMON.

---

# SUPPLEMENT TO THE REPORT

OF THE

## COMMITTEE FOR SCIENTIFIC INQUIRIES.

---

### No. I.

THE following is the letter circulated among Members of the Medical Profession by the General Board of Health in September 1854 :—

*Letter addressed by the President of the General Board of Health to Medical Practitioners.*

SIR,  
General Board of Health, Whitehall,  
September 1854.

MY experience of this Department, brief as it is, has strongly impressed me with a sense of the great want that is now felt of some systematic record of cases of choleraic disease, their treatment, and results, with a view to determine, in so far as may be possible, the best mode of meeting this formidable epidemic.

Hitherto no successful attempt has been made to collect such a record ; and as I find that my feeling of the want of it is very generally shared by the medical profession, I have obtained the sanction of Her Majesty's Government to the nomination of a Medical Council, representing all branches of the profession, and consisting of Dr. Paris, Sir James Clark, Dr. Alderson, Dr. Babington, Dr. Tweedie, Dr. Baly, Mr. Lawrence, Mr. Simon, Professor Owen, Mr. W. B. Ward, Mr. John Bacot, and Dr. Farr.

Under the sanction of this Council the following form of return has been prepared for transmission to all qualified practitioners in the metropolis, to be filled up by them, with a view to obtain their experience of the present epidemic.

The deaths from cholera, as well as from all other causes, are registered in England ; but it is evidently desirable that in this as in other countries the cases of recovery, as well as of death, should be systematically observed and recorded. But this can only be accomplished by the cordial co-operation of all the medical men in practice, which the Board hopes, in this important matter, to obtain, by acting on the advice of a Council in which all branches of the medical profession are represented

By means of a return, in the accompanying forms, the observations of all qualified practitioners on the cases that come under their care may be collected, and made available for determining the laws which regulate choleraic disease, and the effects of the different systems of treatment now in use.

Any return that you may make will be considered strictly confidential, but your name will be recorded as a contributor to the observations, when the general results are made known.

Under the urgent pressure of practice, to keep any accurate record of the cases will often be attended with difficulty; but this difficulty will, I feel assured, be cheerfully encountered by you if you agree with me in thinking a record of the experience of the present generation of medical men as to this epidemic likely to contribute to the alleviation of the sufferings of mankind for the future.

The return on the other side is sent filled up, as a specimen, with blank forms. Any number of such forms you may require will be furnished by this office.

It is recommended that the forms should be filled up from day to day, and returned to the Board, on or before the 15th November next, addressed to the Secretary of the General Board of Health, and the postage of the letter enclosing the return need not be prepaid.

I have the honour to remain, &c.

B. HALL.

#### NOTE.

THE foregoing letter was sent by the General Board of Health to all practitioners whose names appear in the "Medical Directory" of the year 1854.

In pursuance of the promise therein contained, the names of all who contributed returns are now recorded.

Among the names occur some of homœopathic practitioners, from whom returns were received; but the Committee for Scientific Inquiries desire it to be understood that none of these communications have been used in the construction of their Report.



**List of legally-qualified Medical Practitioners, Medical Officers of Hospitals, &c., in England, Scotland, and Wales, who have furnished Returns of Cases of Cholera and Diarrhœa to the General Board of Health.**

(See the President's Letter, p. 67.)

Name.	Address.
Anderson, John, M.R.C.S. & L.S.A.	4, Bedford Terrace, Clapham Rise.
Air, Alex. Cummings, M.R.C.S. -	15, Studley Terrace, Stockwell.
Alexander, Wm., M.D. - -	Dundonald, by Kilmarnock, N.B.
Aspray, Thomas, L.S.A. - -	12, Wilmington Square.
Adcock, Christr., L.S.A. - -	8, Great Charlotte Street, Black-
	friars Road.
Alford, S. S., M.R.C.S. & L.S.A. -	7, Powis Place, Haverstock Hill.
Allen, A., M.R.C.S. - - -	18, Hunter Street, Brunswick Sq.
Arden, H. A., M.R.C.S. & L.S.A.	High Street, Dorchester.
Appleton, H., M.D. Aberdeen,	Mare Street, Hackney.
M.R.C.S. & L.S.A.	
Ayres, Philip B., M.D. Lond., &	6, Upper Portland Place, Wands-
M.R.C.S.	worth Road.
Aitkin, James, M.R.C.S. - -	Ceres, N.B.
Allen, Geo., L.S.A. - - -	14 a, Dean Street, Soho.
Aldis, C. J. B., M.D. - - -	1, Chester Terrace, Pimlico.
Angers, V. P. de Bois, M.D.	2, Salamanca Cottages, Upper
Heidelberg, M.R.C.S.	Church Street, Queen's Elm.
Armstrong, Thos., M.R.C.S. &	Brecon, S.W.
L.S.A.	
Baines, Matthew, M.D. Lond. -	54, Thurloe Square, Kensington.
Burroughs, J. T. R., M.R.C.S. -	3, Lee Place, Lee.
Bunnett, H. B., M.R.C.S. - -	5, Carlton Hill East, St. John's
	Wood.
Bowling, John, M.R.C.S. - -	Pingsworth House, Hammersmith.
Barnwell, Wm., M.R.C.S. & L.S.A.	7, Queen Charlotte Row, New Road.
Barnwell, W. H., M.R.C.S. -	Ditto ditto.
Bainbridge, J. N., M.D. St. And.,	86, St. Martin's Lane.
F.R.C.S. & L.S.A.	
Bainbridge, Wm., M.R.C.S. &	17, Upper Eaton Street, Pimlico.
L.S.A.	
Balding, D. B., M.R.C.S. & L.S.A.	St. Mary-le-bone Infirmary.
Barringer, T. S., M.R.C.S. & L.S.A.	96, St. John Street, Clerkenwell.
Brodrigg, W. P., M.R.C.S. & M.S.A.	9, Bloomsbury Square.
Bennett, J. M., M.R.C.S. - -	46, Upper Baker Street.
Bradley, R. H., M.R.C.S. & L.S.A.	Trafalgar Road, Greenwich.
Bibby, Sam. H., M.R.C.S. & L.S.A.	9, North Audley Street.
Bowra, H. G., M.R.C.S. & L.S.A. -	Brooke Lodge, De Beauvoir Road,
	Kingsland.
Bryant, John, M.D. Ellangen -	Colebrooke Villa, Finchley.

Name.	Address.
Besemerer, Wm., M.R.C.S. & L.S.A.	13, Marlborough Place, Old Kent Road.
Balfour, G. W., M.D. - - -	Westfield, Cramond, N.B.
Bradshaw, H., M.R.C.S. - - -	1, St. James's Road, Liverpool.
Buchanan, G. A., M.R.C.S. - - -	11, Middleton Square, Clerkenwell.
Bonney, F.A.B., L.R.C.S. & L.S.A.	3, Trevor Terrace, Knightsbridge.
Beale, John Sam., M.R.C.S. - - -	87, Harrow Road.
Bullock, Henry, M.R.C.S. & L.S.A.	15, Cumberland Street, Bryanston Square.
Bianchi, Robt., M.R.C.S. & L.S.A.	8, Great Surrey Street, Blackfriars.
Brent, Robt., M.R.C.S. - - -	Sydney Cottage, Woodbury, near Exeter.
Blackmore, J. T., L.S.A. - - -	24, Old Street Road.
Bossy, A. H., L.S.A. - - -	Nelson Terrace, Stoke Newington.
Blundell, John, M.R.C.S. & L.S.A.	St. Helen's, Lancaster.
Clarke, John, M.D. - - -	42, Hertford Street, Mayfair.
Campbell, Wm., M.R.C.S. - - -	New Street, Kilbarchan, N.B.
Christian, J. Stanley, M.D. - - -	1, Ovington Terrace, Brompton.
Carter, Wm. G., M.R.C.S. - - -	1, Manchester Terrace, Kilburn.
Cheyne, R. R., F.R.C.S. & L.S.A.	43, Berners Street.
Clifton, Nath. H., M.R.C.S. - - -	38, Cross Street, Islington.
Covey, W. H., F.R.C.S. - - -	23, Wilton Street, Grosvenor Sq.
Chepmell, E. C., M.D. Edin. - - -	22, Henrietta Street, Cavendish Sq.
Corner, Mat., M.D., M.R.C.S. & L.S.A.	40, Colet Place, Commercial Road.
Cargow, Chas., M.R.C.S. - - -	Dalrymple, Girvan, Ayr, N.B.
Cresswell, John, M.R.C.S. & L.S.A.	Winchmore Hill.
Collambell, Charles, F.R.C.S. & L.S.A.	15, Lambeth Terrace, Lambeth.
Currie, Alex., M.D. - - -	Bowmore, Islay, N.B.
Currie, J. T., L.F.P. & S.G. - - -	10, Norfolk Street, Fitzroy Square.
Clark, Chas., M.R.C.S. - - -	Adelaide Terrace, Notting Hill.
Cameron, Angus, M.D. & L.R.C.S.	Balfon, by Glasgow, N.B.
Cox, Harry, M.R.C.S. & L.S.A. - - -	23, Hill Street, Camberwell.
Cooper, J. P., M.R.C.S. & L.S.A. - - -	Bow Road.
Cunningham, W. J. M., L.S.A. - - -	Hoo, Kent.
Catherwood, Alfred, M.D. Glas. & L.S.A.	21, Charles Square, Hoxton.
Corrie, J., M.D., M.R.C.S. & L.S.A.	Finchley.
Capes, G.A. H., M.R.C.S. & L.S.A.	22, Philpot Lane.
Cowan, Jno. B., M.D. - - -	West Regent Street, Glasgow, N.B.
Collinson, Alfred, M.R.C.S. - - -	81, Connaught Terrace, Paddington.
Cumming, W. S., M.R.C.S. & L.S.A.	2, Vittoria Place, Limehouse.
Christophers, J. C., M.R.C.S. - - -	38, York Street, Portman Square.
Challice, John, M.D. Aberd. & L.S.A.	40, Grange Road, Bermondsey.

Name.	Address.
Chesterman, S., M.R.C.S. & L.S.A.	Banbury.
Colgate, Robt., M.R.C.S. & L.S.A.	Eastbourne, Sussex.
Dale, G. Cornelius, M.R.C.S. & L.S.A.	Commercial Road, St. George's East.
Davis, Robt., M.D., M.R.C.S. & L.S.A.	2, Brunswick Place, City Road.
Davis, H. P., M.R.C.S. & L.S.A.	71, Oakley Square, St. Pancras.
Dickman, David, L.R.C.S.	14, Porchester Place.
Druitt, Robt., M.D. - - -	39a, Curzon Street, Mayfair.
Davis, E., M.R.C.S. & L.S.A. -	17, Upper Belgrave Place, Fimlico.
Davidson, N., L.S.A. - - -	15, Charles Street, Manchester Sq.
Dean, J. H., M.R.C.S. - - -	32, Lucas Street, Commercial Road.
Duncan, Jas., M.B. & M.R.C.S. -	18, Henrietta Street, Covent Garden.
Dickson, John, M.R.C.S. - - -	High Street, Bathgate, N.B.
Duchesne, Clarke, M.R.C.S. & L.S.A.	22, White Lion Street, Norton Folgate.
Devenish, S. W., M.B., M.R.C.S., & L.S.A.	7, Billiter Square.
Dixon, John, M.D., M.R.C.S., & L.S.A.	20, Prospect Row, Bermondsey.
Dickson, James, M.R.C.S. - - -	Roberton-by-Abington, Lanark, N.B.
Dodsworth, F. C., M.R.C.S. - -	Turnham Green.
Dickinson, Thos., M.R.C.S. & L.S.A.	37, Sloane Street, Chelsea.
Douglas, Joseph, M.R.C.S. - - -	1, Sumner Terrace, Brompton.
Dale, Edmund, M.D. St. And., L.S.A.	23, Holborn Hill.
Dupasquier, C. J., M.R.C.S. & M.S.A.	62, Pall Mall.
De Meric, Victor, M.R.C.S. - - -	17, Brook Street, Grosvenor Sq.
Down, Geo., M.R.C.S. & L.S.A. -	4, Upwell Place, Camberwell New Road.
Davies, John, M.R.C.S. & L.S.A.	Bulwark, St. Mary's Brecon.
English, Thomas, M.D. St. And., M.R.C.S. & L.S.A.	32, Upper East Smithfield.
Ellis, H. W. T., M.R.C.S. - - -	Crowle, Lincolnshire.
Edwardes, David, L.S.A. - - -	13, Great Russell Street, Covent Garden.
Engall, Thos., M.R.C.S. - - -	15, Euston Square.
Eve, Richd. Wafford, M.B. & L.S.A.	7, Slade's Place, High Street, Deptford.
Ellis, Cydwelyn, M.D. St. And., M.R.C.S. & L.S.A.	31, Howland Street, Fitzroy Square.
Edmonds, J. W., L.S.A. - - -	Montague Cottage, Southampton Street, Camberwell.
Fowler, Robt., M.D. Edinburgh, M.R.C.S. & L.S.A.	145, Bishopsgate Street Without.

Name.	Address.
Forman, John, M.R.C.S. - -	Fettercairn, N.B.
Forbes, J. G., F.R.C.S. & L.S.A.	9, Devonport Street, Paddington.
Faulker, Wm., M.R.C.S. - -	151, Drury Lane.
Fuller, Hy. Wm., M.D. - -	13, Manchester Square.
Ferguson, Jno., M.R.C.S. - -	Cove-Nigg, N.B.
Farrants, R. J., F.R.C.S. & L.S.A.	5, Regent Terrace, City Road.
Foitherby, H. I., M.B., M.R.C.S., & L.S.A.	40, Trinity Square, Tower Hill.
Gibb, Geo. D., M.D. Montreal, & L.R.C.S.I.	59, Guildford Street, Russell Square.
Giles, Wm., M.R.C.S. - -	1, Canton Place, East India Dock Road, Limehouse.
Grant, Thos. Walker, M.D. Aber., M.R.C.S. & L.S.A.	9, Connaught Terrace.
Gibson, J. B., M.R.C.S. & L.S.A.	3, Spring Place, Bagnigge Wells Road.
Griffith, Samuel Hallett, M.R.C.S. & L.S.A.	2, Charterhouse Square.
Grimbly, Richard, M.R.C.S. & L.S.A.	South Bar, Banbury.
Goodworth, W. H., L.S.A. - -	Hatfield, Yorkshire.
Gibson, J. R., M.R.C.S. & L.S.A. -	115, Holborn Hill.
Grant, Nathl., M.D. Edinburgh, M.R.C.S.	48, York Street, Portman Square.
Graham, Thomas, M.R.C.S. & L.S.A.	2, Oxford Terrace, Upper Holloway.
Goodday, Horatio, M.R.C.S. & L.S.A.	48, Baker Street, Portman Square.
Graham, John, M.R.C.S. - -	Stanley, Perth, N.B.
Goddard, Saml., M.R.C.S. - -	Swan Square, Burslem.
Greenwood, Thos. B., L.S.A. - -	1, Horsleydown Lane, Southwark.
Horton, Fitzwilliam, L.S.A. - -	Glo'ster Lodge, North End, Ful- ham.
Halford, Edw., M.R.C.S. & L.S.A.	The Grove, Hammersmith.
Hart, W. B., M.R.C.S. & L.S.A. -	15, Union Street East, Spitalfields.
Harper, James P., M.D. - -	53, Constitution Street, Leith, N.B.
Halley, Alexr., M.D. - -	14, Queen Anne Street, Cavendish Square.
Hadaway, James, M.R.C.S. & L.S.A.	36, Berwick Street.
Hammond, H. S., M.R.C.S.	Church Street, Edmonton.
Hunter, John, M.R.C.S. - -	115, Upper Street, Islington.
Hall, D. J., M.D., Edin. - -	South Street, Eastbourne, Sussex.
Haffenden, Thos., M.R.C.S. - -	Hanwell.
Harris, H. C., F.R.C.S. & L.S.A. -	1, Windsor Place, City Road.
Harper, Robert, M.R.C.S. & L.S.A.	7, Westbourne Place, Paddington.
Hawkins, James, M.R.C.S. - -	36, Colet Place, Commercial Road.



Name.	Address.
Hicks, J. B., M.D., M.R.C.S., & L.S.A.	High Road, Tottenham.
Haycock, Edwd., M.R.C.S. & L.S.A.	25, Gretton Terrace, Green Street, Bethnal Green.
Hunter, Jas., F.R.C.S. - -	115, Upper Street, Islington.
Humble, W. E., M.D. Lond. - -	66, London Road, Southwark.
Halford, Edwd., M.R.C.S. & L.S.A.	1, City Road.
Harle, Ezra, L.S.A. - - -	48, Cross Street, Islington.
Hensley, F. J., M.D. London, M.R.C.S. & L.S.A. - - -	30, Montague Place, Russell Square.
Hughes, H. W., M.R.C.S. - -	138, Fenchurch Street.
Hunt, Joseph, M.R.C.S. - -	82, Gray's Inn Lane.
Hicks, E. J., L.S.A. - - -	Havant, Hants.
Hall, Wm., M.R.C.S. & L.S.A. -	High Street, Tottenham.
Harling, R. D., M.D. London -	9, Upper Seymour Street, Portman Square.
Harness, H., M.R.C.S. - -	1, Kimbolton Place, Fulham Road.
Hull, Geo., M.D. - - -	1, St. Mary Abbot's Terrace, Kensington.
Hering, Wm., L.S.A. - -	38, Mortimer Street, Cavendish Sq.
Harrison, John, F.R.C.S. - -	Abany, Piccadilly.
Harris, W., F.R.C.S. - - -	16, Dorset Terrace, Clapham Road.
Hutchinson, Jno., M.R.C.S. & L.S.A.	184, Blackfriars Road.
Hudson, Jno., M.D. & M.R.C.S. -	11, Cork Street, Burlington Gardens.
Ingram, Chas., M.R.C.S. & L.S.A.	White Mill, Blandford.
Jefferys, Wm. Edw., M.R.C.S. -	2, St. Augustine's Road, St. Pancras.
Jones, J., M.R.C.S. - - -	5, Seymour Place, Wandsworth Road.
James, W. Reynolds, M.R.C.S. & L.S.A.	68, Milton Street, Dorset Square.
Joseph, Edwd., M.R.C.S. & L.S.A.	15, Great Marylebone Street.
Jones, Jno. Dalston, M.D. Aber., & M.R.C.S.	Queen's Road, Dalston.
Jeanneret, Hy., M.D. Edinburgh, L.R.C.S. & L.S.A.	15, Upper George St., Marylebone.
Jones, Wm., M.R.C.S. - -	51, Strand.
Illingworth, Hy. S., F.R.C.S. & L.S.A.	1, Arlington Street, Piccadilly.
Jones, Walter, M.R.C.S. - -	27, Fetter Lane, Fleet Street.
Jeffery, G. A., M.R.C.S. & L.S.A.	Trinity House, Eastbourne, Sussex.
Kingdon, J. A., M.R.C.S. - -	2, Bank Buildings, Lothbury.
Kesteven, W. B., M.R.C.S. - -	1, Manor Road, Upper Holloway.
Kellock, W. B., M.R.C.S. - -	Stamford Hill.
Knott, Jas. P., L.S.A. - -	Grafton Villa, Blisworth, Northampton.

Name.	Address.
Kirk, J. B., M.D. - - -	Bathgate, Linlithgow, N.B.
Kenny, Josh., M.B. & L.S.A. -	High Street, Stoke Newington.
Knaggs, Robt., M.R.C.S. & L.S.A.	Swindon.
Lovett, Saml., M.R.C.S. & L.S.A.	23, Clare Street, Clare Market.
Lewis, Waller, M.B. Cantab. -	56, Gower Street, Bedford Square.
Leggatt, Richard S., M.R.C.S. & L.S.A.	Eastry, near Sandwich.
Lloyd, Thomas, F.R.C.S. - -	5, New Basinghall Street.
Lyell, John, M.D. - - -	Main Street, Newburgh, Fifeshire, N. B.
Littlewood, J. J., M.R.C.S. & L.S.A.	2, Stonegate Street, Thorne.
Lord, C. F. J., M.R.C.S. - -	Hampstead.
Leese, R. V., M.R.C.S. & L.S.A. -	Norwood.
Littler, Jno. W., M.R.C.S. & L.S.A.	Lower Edmonton.
Leggatt, Alfred, M.R.C.S. & L.S.A.	13, William Street, Lowndes Sq.
Leigh, Wm. Osborne, L.S.A. -	High Street, Deptford.
Lyddon, Jas., M.R.C.S. & L.S.A. -	Okehampton, St. Thomas, Exeter.
Land, W. H., M.R.C.S. & L.S.A. -	Strand Street, Exmouth.
Leonard, Jas., M.R.C.S. & L.S.A. -	2, Salisbury Street, Strand.
Little, W. J., M.D. - - -	34, Brook Street, Havover Square.
Langley, John, L.S.A. - - -	48, Upper Albany St., Regent's Park.
Lane, W. R. F., M.R.C.S. & L.S.A.	2, North Side, Bethnal Green.
Mitchell, Thos. H., M.R.C.S. & L.S.A.	2, Francis Place, Holloway.
Morris, Jas., M.B. Lond. & F.R.C.S.	79, Park Street, Grosvenor Square.
Mould, J. T., M.R.C.S. & L.S.A. -	1, Onslow Crescent, Brompton.
Moss, Edwin, M.R.C.S. - - -	71, Wimpole Street.
Meates, W. C., M.R.C.S. - - -	41, Chester Square.
Miles, Chas., M.R.C.S. & L.S.A. -	13, Conduit St., West, Hyde Park.
M'Donald, Wm., M.R.C.S. - - -	Muirhead Cottage, Cadder, N.B.
Mathew, Jas. Edwd., M.R.C.S. & L.S.A.	54, Conduit Street, Hanover Sq.
Mackie, Archibald, M.D. - - -	Cupar, Fife, N. B.
Macfarlane, D., M.D. - - -	Drymen, Stirlingshire, N. B.
Munday, Chas., M.R.C.S. - - -	86, Snow Hill.
Morgan, John, F.R.C.S. - - -	5, Albion Place, Hyde Park Square.
Meldola, Eleazar, M.D. Giessen, L.S.A.	6, Great Alie Street.
Molloy, Robt., M.D. Aberdeen, M.R.C.S. & L.S.A.	57, Amwell Street, Claremont Square, Clerkenwell.
Malton, Chas. J., F.R.C.S. - - -	6, Stanhope Place, Hyde Park.
Mageniss, Peter, M.R.C.S. - - -	18, Hamilton Place, New Road.
Mitchelson, G. F., M.D. Aber., M.R.C.S. & L.S.A.	11, Ashley Place, Victoria Street, Westminster.
Mackintosh, James Innis, M.D., Edin., L.R.C.S.	St. Olaves Infirmary.

Name.	Address.
McWilliam, J. O., M.D. Edin. -	14, Trinity Square, Tower Hill.
Matheson, L. M., M.R.C.S. -	Portree, Isle of Skye, N.B.
Madden, Jas. M., L.R.C.S. Dub.	Heavitree.
Merry, Wm. H., M.R.C.S. & L.S.A.	Broad Clyst, Exeter.
Mactavish, Colin, M.R.C.S. -	Springbank, Islay, N.B.
Nind, Chas., M.R.C.S. & L.S.A. -	13, Bath Place, Peckham.
Nix, Wm., L.S.A. - - -	11, Savile Row, Mile End Road.
Owen, Harvey K., M.R.C.S.	4, Binfield Place, Clapham Road.
Oliver, Jno. Robt., M.D. - -	28, Harleyford Place, Kennington.
Orr, R. S., M.D. - - -	George Terrace, Dunoon, Argyle-shire, N.B.
O'Shea, Michael Keating, M.R.C.S.	17, Mount Street, Lambeth.
Payne, G. B., M.D. Aber., L.S.A.	58, Denbigh Street, Warwick Square, Pimlico.
Paton James, M.D. - - -	Orr Square, Paisley, N.B.
Palmer, Thomas, M.B., M.R.C.S. & L.S.A. - - - - -	16, Upper Southwick Street, Hyde Park.
Pitman, Henry A., M.D. - -	Montague Place.
Pollock, Robt. Jas., F.R.C.S. & L.S.A.	7, Bath Place, Kensington.
Pullan, Richd., M.R.C.S. - -	Epworth, Bawtry.
Patison, H. W., M.R.C.S. - -	Hirst Street, Renfrew.
Pottle, J. R., L.S.A. - - -	20, Bath Street, St. Luke's.
Parratt, Jas., M.D. St. And., M.R.C.S. & L.S.A.	18, Mount Street, Grosvenor Sq.
Pascall, W. L., L.S.A. - -	17, Charlotte Street, Fitzroy Sq.
Parkes, W. B., M.D. Erlang., F.R.C.S. & L.S.A.	31, Great Marlborough Street.
Pretty, Jno. Rowlison, M.D. St. And., M.R.C.S., & L.S.A.	31, Bayham Terrace, Camden Town.
Paul, Jas Thos., L.S.A. - -	26, Burton Crescent.
Pinder, Edwd., L.R.C.S. & L.S.A.	9, Oakley Terrace, Old Kent Road.
Powell, W. M., M.R.C.S. - -	14, Springfield Place, Wandsworth Road.
Pridham, T. L., M.R.C.S. & L.S.C.	Bideford, Devon.
Pretty, Wm., M.R.C.S. - -	70, Mornington Road.
Pullin, Thos. Hy. S., M.R.C.S. -	Sidmouth, Devon.
Paine, H. J., M.R.C.S. - -	Cardiff. -
Rogers, W. R., M.D., Heidelberg, L.S.A.	56, Berners Street, Marylebone.
Robarts, H. P., F.R.C.S. - -	11, Great Coram Street, Russell Sq.
Rushforth, Richard, M.D. Erlang., L.S.A.	44, Upper Norton Street, Marylebone.
Robinson, Thomas, M.R.C.S. & L.S.A.	Southampton Street, Pentonville.
Radcliffe, Derwentwater, M.R.C.S. & L.S.A.	51, Strand.



Name.	Address.
Ridge, Benjn., M.D., M.R.C.S., & L.S.A.	High Street, Putney.
Roods, H. C., M.D. & M.R.C.S. -	32, Bloomsbury Street, Bedford Square.
Ruttledge, Jas., M.D. Glasgow -	19, George Street, Hanover Square.
Rose, John, M.R.C.S. & L.S.A. -	1, Trinidad Place, Islington.
Robertson, John, L.R.C.S. -	Stanley, Perthshire, N. B.
Riches, Thos. H., L.S.A. - -	London Street, Greenwich.
Robins, Wm., L.S.A. - -	41, Oxford Terrace, Hyde Park.
Roper, George, M.R.C.S. & L.S.A.	180, High Street, Shoreditch.
Read, Reginald, L.S.A. - -	1, Guildford Place, Russell Square.
Read, Septimus, L.S.A. - -	41, Jewin Street, Cripplegate.
Sanderson, A. M., F.R.C.S. -	19, Bridge Street, Musselburgh, N. B.
Smith, Anderson, M.R.C.S. & L.S.A.	4, Northumberland Terrace, Bagnigge Wells Road.
Stewart, Hy. C., M.R.C.S. -	42, Grove Road, St John's Wood.
Stevens, Wm., L.S.A. - -	6, Lansdowne Villas, Fulham Road.
Snell, Edmund, M.R.C.S. - -	18, Crown Row, Mile End Road.
Smith, Geo., M.R.C.S. - -	Axbridge.
Sewell, Eade, M.R.C.S. & L.S.A.	Englefield Road, Hackney.
Stokes, Hy. J., M.D. Edin. -	23, Portland Place, Islington.
Semple, Wm., M.R.C.S. & L.S.A.	38, Upper Street, Islington.
Scatliff, Arthur, M.R.C.S. & L.S.A.	1, Grosvenor Place, Brixton.
Schulhof, Maurice, M.D. - -	7, Suffolk Place, Pall Mall, East.
Statham, Hugh, M.R.C.S. & L.S.A.	16, Belmont Place, Wandsworth Road.
Stevens, N. Hy., M.R.C.S. - -	2, Finsbury Place South, Finsbury Square.
Smiles, John Finch, L.S.A. -	South Bar Street, Banbury.
Simmonds, H. M., M.R.C.S. & L.S.A.	76, Albany Road, Camberwell.
Smyth, Jno. Edwd., M.R.C.S. & L.S.A.	2, China Terrace, Lambeth.
Smith, E. Pye, F.R.C.S.E. - -	Mare Street, Hackney.
Sandys, Saml., M.R.C.S. & L.S.A.	4, Francis Terrace, Kentish Town.
Sutherin, Hy., M.R.C.S. - -	22, Church Row, Pancras Road.
Scarnell, S. F., L.S.A. - -	1, Langton Place, Vassall Road, Brixton.
Spooner, Edwd. O., F.R.C.S. -	White Mill, Blandford.
Smith, Chas., M.R.C.S. & L.S.A.	1, Upper Craven Place, Kentish Town.
Spettigue, John, M.R.C.S. & L.S.A.	3, Bieton Terrace, Littleham, Exmouth.
Smith, Thomas Hy., M.R.C.S. & L.S.A.	13, John Street, Berkeley Square.
Thompson, J. B., L.R.C.S. - -	Alva, N. B.



Name.	Address.
Turner, Saml. Mayer, F.R.C.S. -	Newcastle-under-Lyne, Staffordshire.
Taylor, John, M.R.C.S. & L.S.A.	64, Queen's Road, Bayswater.
Tweed, J. J., M.R.C.S. & L.S.A. -	14, Upper Brook Street, Grosvenor Square.
Trousdale, Alfred, M.R.C.S. -	Epworth near Bawtry.
Tweedie, A. C., M.R.C.S. - -	Galashiels, N. B.
Tapson, John, M.D., M.R.C.S. & L.S.A.	High Street, Clapham.
Timms, Godwin Wm., M.D. Lond., & M.R.C.S.	28, Paddington Green.
Townley, Jas., F.R.C.S. & L.S.A.	Kennington Common.
Thomas, Jno. Hy., M.R.C.S. & L.S.A.	Upper Street, Islington.
Tanner, Robt, M.R.C.S. & L.S.A.	56, Manchester Street, Manchester Square.
Turnbull, William, L.S.A. - -	62, Great Russell St., Bloomsbury.
Todd, J. M., M.R.C.S. & L.S.A. -	10, Kent Terrace, New Cross Road, Deptford.
Taylor, Thos., M.R.C.S. - -	High Street, Cricklade.
Vinen, J. Northcote, L.S.A. -	114, Tooley Street.
Ward, N. B., M.R.C.S. - -	14, Clapham Rise.
Wetherfield, John, M.R.C.S. -	1, Henrietta Street, Covent Garden.
Wallace, Richd., F.R.C.S. & L.S.A.	1, Trafalgar Place, West Hackney Road.
Woodforde, W. T. G., M.D. Lond.	37, High Street, Bow.
Whittle, E., M.D. Lond. - -	129, Mill Street, Liverpool.
Watson, John, L.S.A. - -	13, Southampton Street, Bloomsbury.
Webb, F. C., M.D. Edin., M.R.C.S.	39, Great Coram Street.
Ward, Martindale, M.R.C.S. & L.S.A.	1, King's Parade, Chelsea.
Weston, E. J., M.R.C.S. - -	17, St. Mark's Crescent, Regent's Park.
Wise, R. S., M.D., M.R.C.S. -	High Street, Banbury.
Wright, Edwd., M.R.C.S. & M.S.A.	5, Kennington Row, Kennington.
Weathers, George, M.R.C.S. & L.S.A.	41, High Street, Camden Town.
Waggett, Jno., M.D. - -	1, Norland Square, Notting Hill.
Walker, David, R. G., M.R.C.S. -	Budleigh, Salterton, Devon.
Williamson, Thos., M.D. - -	33, Charlotte Street, South Leith, N. B.
Winnard, J. T., M.R.C.S. - -	Standish-gate, Wigan.
Young, Chas., M.R.C.S. & L.S.A.	Turner's Hill, Worth, Sussex.
Young, Jas. Forbes, M.D. Edin., & L.S.A.,	17, Upper Kennington Lane.

## HOSPITALS, &amp;c.

Name.	Address.
<i>St. Mary's Hospital.</i>	
Alderson, Jas., M.D. - -	20, Berkeley Square.
Chambers, Thos. K., M.D. - -	1, Hill Street, Berkeley Square.
Jones, C. Handfield, M.D. - -	1, Southwick Place, Hyde Park.
Sibson, Francis, M.D. - -	40, Brook Street, Grosvenor Square.
Sieveking, E. H., M.D. - -	3, Bentinck Street, Manchester Square.
<i>St. Thomas's Hospital.</i>	
Barker, T. A., M.D. - -	71, Lower Grosvenor Street.
Whitfield, Richd. G. - -	Resident Medical Officer at St. Thomas's Hospital.
<i>Chelsea Hospital.</i>	
Maclachlan, D., M.D. - -	Royal Hospital, Chelsea.
<i>St. George's Hospital.</i>	
Barclay, A. Whyte, M.D. - -	Medical Registrar to the Hospital. 23a, Bruton Street, Berkeley Sq.
<i>Westminster Hospital.</i>	
Fincham, George T., M.D. - -	28, Chapel Street, Belgrave Square.
Roe, Hamilton, M.D. - -	26, Upper Brook Street, Grosvenor Square.
<i>Charing Cross Hospital.</i>	
Golding, B., M.D. - -	King William Street, Strand.
<i>Middlesex Hospital.</i>	
Sibley, Septimus W. - -	Medical Registrar to the Hospital.
<i>University College Hospital.</i>	
Hillier, Thos., M.B., London -	Resident Medical Officer.
<i>Royal Free Hospital, Gray's Inn Lane.</i>	
Curgenven, J. Brendon - -	Resident Medical Officer.
<i>St. Bartholomew's Hospital.</i>	
Martin, Robert, M.D. - -	St. Bartholomew's Hospital.
<i>Bethnal House Lunatic Asylum.</i>	
Austin, Thos., M.R.C.S. & L.S.A.	Bethnal House, Bethnal Green.
<i>London Hospital.</i>	
Fraser, Patrick, M.D. - -	62, Guildford Street, Russell Square.
Davies, Herbert, M.D. - -	23, Finsbury Square.

Name.	Address.
<i>Seamen's Hospital</i> ("Dreadnought").	
Complin, Edwd. Jno., M.R.C.S. & L.S.A.	Hospital Ship "Dreadnought."
<i>Royal Hospital, Greenwich.</i>	
Liddell, Sir Jno., M.D.	
M'Ternan, Jas.	
Nesbet, Alex., M.D.	
Whitmarsh, John.	
<i>Christ's Hospital, Newgate Street.</i>	
Stone, Thos., M.R.C.S. - - -	Christ's Hospital.
<i>H.M. Dockyard, Woolwich.</i>	
Browning, Colin Arrott, M.D. -	Surgeon to H.M. Dockyard, Woolwich.
<i>Middlesex House of Detention.</i>	
Wakefield, Henry, M.R.C.S. -	52, Russell Square.
<i>Caledonian Asylum.</i>	
Bradley, C. L., M.R.C.S. - -	4, Belitha Villas West, Barnsbury Park.
<i>Royal Military Asylum.</i>	
Balfour, T. G., M.D. - - -	Surgeon to Royal Military Asylum.
<i>Naval Hospital, Walmer.</i>	
Johnston, J. W., M.D. - - -	Deputy Inspector of Naval Hospitals.
<i>Homœopathic Hospital.</i>	
Yeldham, Stephen, M.R.C.S.E. & L.S.A.	
Leadham, Thos. R., M.R.C.S.E. & L.S.A.	
Mackern, Thomas, L.R.C.S.I.	
Cameron, Hugh, M.R.C.S.E.	
Wielobycki, S., M.D.	
Hill, John Hutton, M.D.	
Wyld, Geo., M.D.	
Baikie, Robt., M.D.	
Henry, Alexr., M.D.	
Smith, Dan., M.R.C.S.E. & L.S.A.	
Morgan, William, M.R.C.S.E.	
Quin, F. P., M.D.	
Hamilton, Edward, M.D.	
	See Parliamentary Return, Cholera, No. 255, Session 1855.

## Instructions and Forms for Returns of Cases.

### A.—DIARRHŒA.

#### SHEET OF INSTRUCTIONS AND EXAMPLES.

(Attention is particularly requested to the following Instructions, as the Value of the Returns will depend on their Uniformity.)

INSTRUCTION I.—The following *degrees or stages of the disease* should, when possible, be distinguished :—

<i>Diarrhœa.</i>	{	1. <i>Simple</i> .—Alvine discharges liquid, but fœcal. Vomiting and cramps absent.
		2. <i>Choleraic</i> .—Alvine discharges very copious, watery, still tinged with bile. Vomiting generally present, but not continued or urgent. Cramps absent.

ALL CASES OF “DIARRHŒA” ARE TO BE ENTERED ON THE SHEET A. AS SOON AS THEY ARE SEEN.

SO SOON AS THE “CHOLERAIC DIARRHŒA” PASSES INTO “CHOLERA,” THE FACT SHOULD BE RECORDED IN THE PROPER COLUMN OF THIS RETURN (A), AND THE PEN SHOULD BE DRAWN THROUGH THE CASE, ALL THE PARTICULARS OF WHICH SHOULD THEN BE TRANSFERRED TO THE “RETURN B.,” AND ITS SUBSEQUENT COURSE BE THERE RECORDED.

INSTRUCTION II. *Absence of Stages*.—The commencement of “Choleraic Diarrhœa” is generally marked by vomiting; and when vomiting exists from the very beginning, the word “*absent*” or “*abs.*” may be written opposite “Simple Diarrhœa” in the Return.

INSTRUCTION III. *Dates*.—When the time of commencement of either stage cannot be ascertained, the words “*not known*” or “*n. k.*” should be written in the place of the date: but the time of commencement may generally be stated approximately. (See Instruction III. Return B.)

INSTRUCTION IV. *Treatment*.—The nature of the treatment should be indicated in the Return as concisely as possible. When any uniform and definite plan of treatment is adopted in a series of cases, that plan should be accurately described in the space for “REMARKS,” and should be indicated in the Table by one or two words, as “*Conf. Arom. c. Op.*,” “*Sulph. Ac.*,” &c.

---



# A.—DIARRHŒA—continued.

## FORM OF RETURNS.

(Attention is particularly requested to the Sheet of Instructions and Examples.)

### GENERAL RETURN OF THE NUMBER OF CASES OF DIARRHŒA NOT PASSING INTO CHOLERA.

Total Number of Cases.	Cases of Simple Diarrhœa not passing into further Stages.	Cases of Choleraic Diarrhœa not passing into further Stages.

### ALL CASES OF DIARRHŒA, WITH PARTICULARS OF THEIR DURATION AND TREATMENT.

No. of Case.	Residence when attacked (Street and number of House.)	Sex.		Age (last Birth-day.)	Rank and Occupation. (If Master, write Master after Name of Occupation.)	Degrees or Stages of the Disease.	Dates of their Commencement.		Date of Termination.			TREATMENT.	
		M.	F.				Day.	Hour.	In Recovery.	In Death.	In Cholera.	When commenced.	Its NATURE.
						Simple Diarrhœa - Choleraic Diarrhœa -						Day.	Hour.
						S. Diarrh. - Ch. Diarrh. -							
						&c. &c.							

#### REMARKS.

The above is a correct Return of Cases of Cholera observed and treated by me.

Name

Professional Titles

Residence

Street

Parish

Registration sub-district

## Instructions and Forms for Returns of Cases.

### B.—CHOLERA.

#### SHEET OF INSTRUCTIONS AND EXAMPLES.

(Attention is particularly requested to the following Instructions, as the Value of the Returns will depend on their Uniformity.)

INSTRUCTION I.—*The following Degrees or Stages of the Disease are generally recognized by the Medical Profession, and wherever it is possible should be distinguished. The terms adopted to designate them are in common use. All the stages are not present in every case.*

- |                  |  |
|------------------|--|
| <i>Diarrhœa.</i> | { 1. <i>Simple</i> .—Alvine discharges frequent and liquid, but fœcal. Vomiting and cramps absent.<br>2. <i>Choleraic</i> .—Alvine discharges very copious, watery, still tinged with bile. Vomiting generally present, but not continued or urgent. Cramps of extremities absent.   |
| <i>Cholera.</i>  | { 3. <i>Without Collapse</i> .—Alvine discharges watery, colourless, with white flakes (rice-water). Vomiting commonly urgent. Cramps of extremities frequent and severe. Eyes somewhat sunken. Temperature of surface lowered. Pulse small and feeble. Urine not secreted.<br>4. <i>With Collapse</i> .—Surface of face and extremities quite cold, often wet. Face and hands much shrunk, and more or less deeply livid. Cramps present. Pulse at wrist absent or scarcely to be felt. Veins of extremities contracted to dark threads. Urine not secreted. Voice usually much altered and feeble.<br>5. <i>Consecutive Fever</i> .—Temperature of surface more or less restored. Pulse distinct, sometimes full and throbbing. Veins more or less filled. Face less shrunk, or even full and deeply flushed. Drowsiness passing into stupor. Alvine discharges again containing bile. Urine, in most cases, still suppressed. |

INSTRUCTION II. *Absence of Stages*.—The absence of any one or more of the earlier stages should be indicated by the word "*absent*" or "*abs.*," written opposite the deficient stage in the place of the date of commencement. The fact of the disease not reaching the later stages will be sufficiently shown by the mode of termination of the case, ("*death*," or "*recovery*,") being written opposite the stage at which the disease ceased.

INSTRUCTION III. *Dates*.—When the time of commencement of a particular stage cannot be ascertained, the words "*not known*," or "*n. k.*," should be written in the place of the date. But although the precise hour of the commencement of each stage cannot be determined exactly, except in rare instances, it may generally be stated approximately by taking some intermediate time between a known period when the symptoms of the particular stage were entirely absent, and one in which they were clearly developed.

INSTRUCTION IV. *Recovery*.—The date of recovery should be fixed at the time when all the symptoms of the disease, and all marked disturbances of health directly resulting from it, have disappeared, although some degree of debility may remain.

INSTRUCTION V. *Treatment*.—The nature of the treatment should be indicated in the Table as concisely as possible. When any uniform and definite plan of treatment is adopted in a series of cases, that plan should be accurately described in the space for "REMARKS," and should be indicated on the Table by one or two words, as "*Salines*," "*Calomel c. Op.*," "*Stimulants*," "*Sulph. Ac.*" &c.

INSTRUCTION VI.—If any patient at the time of the attack was already suffering from another disease, the nature of that disease and the treatment used for it should be mentioned in the "REMARKS."

---

# B.—CHOLERA—continued.

## FORM OF RETURNS.

(Attention is particularly requested to the Sheet of Instructions and Examples.)

### GENERAL RETURN OF THE NUMBER OF CASES OF CHOLERA OBSERVED.

Total Number of Cases.	Cases of Cholera not passing into complete Collapse.	Cases of complete Collapse.	Number of foregoing Cases in which Consecutive Fever supervened.

### ALL CASES OF CHOLERA, WITH PARTICULARS OF THE DURATION AND TREATMENT OF THE SEVERAL STAGES IN EACH CASE.

No. of Case.	Residence when attacked (Street and number of House.)	Sex.		Age (last Birth-day.)	Rank and Occupation. (If Master, write Mast. after the Name of Occupation.)	Degrees or Stages of the Disease.	Dates of their Commencement.		Termination of the Case.		Dates of Death or Recovery.		Treatment in the several Stages.
		M.	F.				Day.	Hour.	Death.	Recovery.	Day.	Hour.	
						Simple Diarrhœa - Choleraic Diarrhœa - Cholera (without collapse) Collapse - Consecutive Fever -							
						S. Diarrh. - Ch. Diarrh. - Cholera - Coll. - Consec. F. -							
						&c. &c.							

#### REMARKS.

The above is a correct Return of Cases of Cholera observed and treated by me.

Name \_\_\_\_\_  
Professional Titles \_\_\_\_\_  
Residence \_\_\_\_\_ Street \_\_\_\_\_ Parish \_\_\_\_\_  
Registration sub-district \_\_\_\_\_



## Memorandum on the foregoing "Instructions and Forms for Returns of Cases of Diarrhœa and Cholera."

IN the Statistical Section of the Committee's Report, it is shown that the degrees of danger to which patients are exposed—or, on the other hand, their chances of recovery—are very different in the different stages of choleraic disease. As long as they suffer only from the diarrhœa stage, their chances of recovery are probably not less than 50 to 1. But on the symptoms of unequivocal cholera supervening, their chances of dying or recovering are nearly equal. If they pass into collapse, the chances are 2 to 1 against their recovery; while, if they survive this stage, but do not at once recover, their chances in the consecutive fever stages are 2 to 1 in favour of their recovery.

The risks attending the different stages of the diarrhœa and cholera had not before been measured. Yet it was well known that the danger increased as the disease advanced to the stage of complete cholera with collapse. The Medical Council, therefore, felt that the relative powers of different modes of treating cholera could not be estimated, unless the relative severity of the cases submitted to the several methods of cure and the stages at which the treatment was commenced in each case were known; and the Forms of Returns issued to medical practitioners (and reprinted in the Appendix) were drawn up with reference to this object. The degrees or stages of the disease which have been generally recognised by experienced writers, viz.:—simple diarrhœa, choleraic diarrhœa, cholera without collapse, cholera with collapse, and the consecutive fever, were distinguished in the Tabular Forms, and were described in brief definitions. Difficulty may sometimes have been experienced in identifying all the phases of particular cases with the stages thus defined; for the different stages, for the most part, pass one into another by almost imperceptible gradations; entire stages may be absent, or may, from their short duration, escape observation; and certain important symptoms may occur earlier or later than the stage of which they are usually characteristic. But in the large majority of cases, it will be possible to determine what is the predominant character of the group of symptoms present at a given time.

The simple diarrhœa does not differ from the ordinary sporadic diarrhœa in any known characters, except it be in the entire or nearly entire absence of pain.

The choleraic diarrhœa is, on the other hand characterized by the extraordinary copiousness of the discharges, and their watery nature, though bile is still present in them, and usually by the early occurrence of vomiting; and, further, by the exhaustion consequent on the loss of fluids, which is both felt by the patient and manifested in his countenance and pulse.

The distinction between choleraic diarrhoea and cholera without collapse will sometimes be attended with difficulty, or at least require care. Slight cramps occur in cases which are shown by all other characters to be as yet only in the stage of choleraic diarrhoea. The discharges may assume very early the rice-water character, while the other symptoms are not proportionally severe. In such cases the consideration of the symptoms collectively, with especial regard to the amount of exhaustion present and the rapidity of its increase, will usually lead to a correct judgment as to the existing degree or stage of the disease.

As the danger in cholera is so much affected by the presence or absence of collapse, it is most important that in all numerical reports of cases intended to illustrate the results of treatment, the character of the cases in this particular should be correctly stated. The essential phenomena of collapse are the more or less diminished volume of the blood and its slow movement, both of which may be best seen in the superficial veins. The complete or nearly complete absence of pulse at the wrist, the loss of heat, and the shrinking of the tissues in some degree are the necessary consequences of this state of the circulating fluid. But the dark or livid colour of the surface of the body is subject to great variety in cases of equally advanced collapse ; it appears to be dependent, at least in great part, on the colour and other conditions of the skin in health, and is imperfectly developed in persons of fair and pale complexion. It is, therefore, not an essential phenomenon of collapse.

The stage denominated consecutive fever cannot be precisely defined. For the morbid conditions which too often succeed to the collapse of cholera are various. Those best known are the imperfect reaction from collapse, the state of uræmia, and the states of sympathetic fever and other disturbance of the general system attending inflammation of the lungs, ulceration and sloughing of the intestines, or suppuration of the parotids. The same mode of treatment cannot be equally applicable to all these different forms of consecutive disease. A valuable addition, therefore, will be made to our knowledge, if in future returns it be stated in each case what was the form of consecutive disease submitted to the treatment.

---

## No. II.

Statistical Tables and other Documents illustrating the Report  
of the Medical Council's Committee for Scientific Purposes.

I.—The FIRST SERIES of TABLES (I. to XI.) was compiled by Mr. Lindsey Blyth, from the Medical Returns (A. B.), which were filled up by the Medical Practitioners of London, and of some other districts in which cholera prevailed.

II.—The SECOND SERIES of Tables (I. to X.) was compiled by Mr. Henry Edwards, assisted by Mr. C. Taylor, and others, from the Registers of Deaths, with the permission of the Registrar-General.

III.—The THIRD SERIES of TABLES, comprising Deductions and Numerical Results, partly taken from the preceding Tables, and partly from the Registrar-General's Weekly Tables. The additional Tables are calculated by Mr. F. J. Williams.

## FIRST SERIES.

TABLE I.

Where the Cases were observed.	CHOLERA.			CHOLERA and DIARRHŒA.	DIARRHŒA.		
	Cases.	Recoveries.	Deaths.	Total Cases, Cholera and Diarrhœa.	Deaths.	Recoveries.	Cases.
In Hospitals - - -	1,502	739	763	2,190	8	680	688
Out of Hospitals - -	1,686	982	704	18,458	101	16,671	16,772
Total in London -	3,188	1,721	1,467	20,648	109	17,351	17,460
Provincial Districts, England	586	348	233	2,762	40	2,136	2,176
Provincial Districts, Scotland	497	254	243	1,162	7	658	665
Total - - -	4,271	2,323	1,948	24,572	156	20,145	20,301

FIRST SERIES—*continued.*

TABLE II.—THREE GROUPS of CASES of DIARRHŒA, distinguishing those that were returned in the prescribed Forms.

—	Cases.	Recoveries.	Deaths.
(1.) Diarrhœa, simple and choleraic, (cases returned only in the aggregate) -	9,103	9,075	28
(2.) Diarrhœa, simple and choleraic, (which have been returned in detail in prescribed forms) - - -	10,323	10,200	123
(3.) Diarrhœa, simple and choleraic, returned in the prescribed forms, (passing into other diseases) - - -	15	9	6
Total - - - -	19,441	19,284	157

None of these cases, even when fatal, passed into the stages of cholera or collapse; fifteen, however, passed into dysentery, fever, and other diseases.

TABLE III.—CASES of CHOLERA, distinguishing their Form at the commencement, as stated in the Returns.

Returned as commencing in	Cases.	Recoveries.	Deaths.
(1.) Diarrhœa - - - -	1,317	699	618
(2.) Choleraic Diarrhœa - - -	360	188	172
(3.) Cholera - - - -	199	86	113
(4.) Collapse - - - -	6	1	5
(5.) Mode of origin unknown - -	1,816	947	869
Total - - - -	3,698	1,921	1,777



FIRST SERIES—*continued.*

TABLE IV.—The same CASES of CHOLERA as in Table III., distinguishing the MODES of TERMINATION.

Returned as terminating in the stage of	Cases.	Recoveries.	Deaths.
(1.) Cholera (without collapse) - -	924	813	111
(2.) Cholera (with collapse) - -	1,798	391	1,407
(3.) Consecutive fever (preceded by cholera or collapse) - - - -	874	625	249
(4.) Stage of termination doubtful - -	102	92	10
Total - - - -	3,698	1,921	1,777

TABLE V.—CASES of CONSECUTIVE FEVER, distinguishing those preceded by COLLAPSE, and those preceded by CHOLERA without COLLAPSE.

—	Total Cases.	Recoveries.	Deaths.
Preceded by cholera - - -	874	625	249
Namely, by cholera without collapse -	223	197	26
By cholera with collapse - - -	585	386	199
And by cholera in which this point was not recorded - - - - }	66	42	24

TABLE VI.—Cases of DIARRHŒA and CHOLERA, distinguishing the Mode of Origin and TERMINATION (including those Cases of Diarrhœa returned in the aggregate, without the details of Age and other circumstances).

		TERMINATING IN															STAGE OF TERMINATION DOUBTFUL.		
		TOTAL.			DIARRHŒA.			CHOLERA.			COLLAPSE.			CONSECUTIVE FEVER.			DYSENTERY AND OTHER DISEASES.		
		Cases.	Recoveries.	Deaths.	Cases.	Recoveries.	Deaths.	Cases.	Recoveries.	Deaths.	Cases.	Recoveries.	Deaths.	Cases.	Recoveries.	Deaths.	Cases.	Recoveries.	Deaths.
		23,139	21,205	1,934	19,426	19,275	151	924	813	111	1,798	391	1,407	874	625	249	15	9	6
		20,171	947	19,426	19,275	151	310	290	20	793	181	612	561	404	157	15	9	6	1
Commencing in diarrhœa		21,118																	
Returned as																			
Commencing in cholera		199	86	113	—	—	22	20	2	118	20	98	59	46	13	—	—	—	—
Commencing in collapse		6	1	5	—	—	—	—	—	5	1	4	1	—	1	—	—	—	—
Mode of origin unknown		1,816	947	869	—	—	592	503	89	882	189	693	253	175	78	—	89	80	9

FIRST SERIES—*continued.*

TABLE VII.—Of the DURATION of CASES of RECOVERY and DEATH from CHOLERA and from DIARRHŒA in

## MALES.

Duration.	CHOLERA.			DIARRHŒA.			Duration.
Days.	Cases.	Recoveries.	Deaths.	Cases.	Recoveries.	Deaths.	Days.
0—	58	15	43	130	127	3	0—
1—	201	17	184				1—
2—	130	7	123				2—
3—	55	1	54				3—
0—	444	40	404	130	127	3	0—
1—	300	77	223	602	594	8	1—
2—	222	117	105	519	516	3	2—
3—	180	127	53	338	329	9	3—
4—	151	110	41	209	205	4	4—
5—	105	84	21	161	155	6	5—
6—	87	69	18	116	112	4	6—
7—	81	64	17	81	77	4	7—
14—	256	215	41	202	197	5	14—
21—	60	54	6	51	51	0	21—
28—	29	27	2	12	10	2	28—
Not known	17	8	9	0	0	0	Not known.
Total -	1932	992	940	2421	2373	48	Total.

TABLE VIII.—Of the DURATION of CASES of RECOVERY and DEATH from CHOLERA and from DIARRHŒA in

## FEMALES.

Duration.	CHOLERA.			DIARRHŒA.			Duration.
Days.	Cases.	Recoveries.	Deaths.	Cases.	Recoveries.	Deaths.	Days.
0—	48	16	32	154	151	3	0—
1—	146	15	131				1—
2—	144	10	134				2—
3—	34	1	33				3—
0—	372	42	330	154	151	3	0—
1—	290	85	205	671	669	2	1—
2—	181	100	81	600	592	8	2—
3—	162	109	53	363	360	3	3—
4—	120	87	33	295	290	5	4—
5—	80	59	21	189	188	1	5—
6—	76	53	23	129	126	3	6—
7—	96	71	25	106	101	5	7—
14—	232	195	37	264	261	7	14—
21—	57	54	3	62	61	1	21—
28—	19	17	2	13	12	1	28—
Not known	18	14	4	0	0	0	Not known.
Total -	1703	806	817	2850	2811	59	Total.

The Table may be read thus: of 1,703 cases of CHOLERA in Females, the duration was not returned in 18 instances; 43 cases terminated in less than 6 hours, or the  $\frac{1}{2}$  of a day; 146 in 6 and less than 12 hours (or  $\frac{1}{2}$  to  $\frac{1}{2}$  day), and so on; thus, 372 cases terminated in less than 24 hours, or in one day, namely, 42 in recovery, 330 in death.

FIRST SERIES—*continued.*

TABLE IX.—CASES of CHOLERA and DIARRHŒA occurring in PERSONS of different Ages, and distinguishing the Number of RECOVERIES and DEATHS at each Age.

Ages.	CHOLERA and DIARRHŒA.			CHOLERA.			DIARRHŒA.		
	Cases.	Recoveries.	Deaths.	Cases.	Recoveries.	Deaths.	Cases.	Recoveries.	Deaths.
0—	196	163	33	23	10	13	173	153	20
1—	192	171	21	22	12	10	170	159	11
2—	187	142	45	68	27	41	119	115	4
3—	194	125	69	98	32	66	96	93	3
4—	170	117	53	83	31	52	87	86	1
	939	718	221	294	112	182	645	606	39
5—	202	137	65	131	67	64	71	70	1
10—	494	379	115	240	128	112	254	251	3
15—	398	330	68	185	120	65	213	210	3
20—	705	597	108	299	195	104	406	402	4
25—	816	692	124	341	218	123	475	474	1
30—	969	821	148	357	210	147	612	611	1
35—	866	698	168	362	196	166	504	502	2
40—	795	648	147	302	158	144	493	490	3
45—	584	455	129	240	112	128	344	343	1
50—	499	392	107	185	80	105	314	312	2
55—	411	305	106	205	100	105	206	205	1
60—	378	282	96	154	65	89	224	217	7
65—	285	213	72	112	46	66	173	167	6
70—	266	186	80	103	28	75	163	158	5
75—	142	105	37	50	15	35	92	90	2
80—	81	53	28	33	7	26	48	46	2
85—	41	27	14	13	3	10	28	24	4
90—	8	6	2	3	1	2	5	5	—
95—	2	2	—	1	1	—	1	1	—
100	1	—	1	1	—	1	—	—	—
N.K.	74	66	8	24	16	8	50	50	—
Total	8,956	7,112	1,844	3,635	1,878	1,757	5,321	5,234	87



FIRST SERIES—*continued.*

TABLE X.—CASES of CHOLERA and DIARRHŒA occurring in MALES of different Ages, and distinguishing the Number of RECOVERIES and DEATHS at each age.

Ages.	CHOLERA and DIARRHŒA.			CHOLERA.			DIARRHŒA.		
	Cases.	Recoveries.	Deaths.	Cases.	Recoveries.	Deaths.	Cases.	Recoveries.	Deaths.
0—	526	394	132	180	70	110	346	324	22
5—	108	68	40	72	33	39	36	35	1
10—	265	207	58	128	72	56	137	135	2
15—	222	182	40	113	76	37	109	106	3
25—	726	602	124	340	219	121	386	383	3
35—	872	696	176	395	221	174	477	475	2
45—	632	489	143	279	136	143	353	353	—
55—	405	303	102	180	79	101	225	224	1
65—	291	212	79	124	51	73	167	161	6
75—	221	158	63	81	21	60	140	137	3
85—	71	42	29	28	4	24	43	38	5
95—	2	2	—	—	—	—	2	2	—
100	—	—	—	—	—	—	—	—	—
N.K.	12	10	2	12	10	2	—	—	—
Total	4,353	3,365	988	1,932	992	940	2,421	2,373	48

TABLE XI.—CASES of CHOLERA and DIARRHŒA occurring in FEMALES of different Ages, and distinguishing the Number of RECOVERIES and DEATHS at each Age.

Ages.	CHOLERA and DIARRHŒA.			CHOLERA.			DIARRHŒA.		
	Cases.	Recoveries.	Deaths.	Cases.	Recoveries.	Deaths.	Cases.	Recoveries.	Deaths.
0—	413	324	89	114	42	72	299	282	17
5—	94	69	25	59	34	25	35	35	—
10—	229	172	57	112	56	56	117	116	1
15—	176	148	28	72	44	28	104	104	—
25—	795	687	108	300	194	106	495	493	2
35—	963	823	140	324	185	139	639	638	1
45—	747	614	133	263	134	129	484	480	4
55—	505	394	111	210	101	109	295	293	2
65—	372	283	89	142	60	82	230	223	7
75—	187	133	54	72	22	50	115	111	4
85—	51	38	13	18	6	12	33	32	1
95—	8	6	2	4	2	2	4	4	—
100—	1	—	1	1	—	1	—	—	—
N.K.	12	6	6	12	6	6	—	—	—
Total	4,553	3,697	856	1,703	886	817	2,850	2,811	39

FIRST SERIES—*continued.*

TABLE XII.—PROPORTIONAL NUMBER of DEATHS, at Twelve different Ages, to every 100 Cases of CHOLERA and DIARRHŒA, in the year 1854.

Age.	To 100 Cases of CHOLERA, at each Age, the Number of Deaths.	To 100 Cases of DIARRHŒA, at each Age, the Number of Deaths.
All Ages	48'3	1'6
0—	61'8	6'0
5—	48'5	1'4
10—	46'7	1'2
15—	34'9	1'4
25—	35'4	0'6
35—	43'4	0'3
45—	50'1	0'5
55—	53'8	0'6
65—	58'2	3'2
75—	71'4	2'7
85—	78'3	0'8
95 & upwards.	60'0	—

The Table may be read thus :—Out of 100 cases of Cholera, which occurred in 1854, at the age of 25 and under 35, there were 35'4 deaths and 64'6 recoveries; out of 100 cases of Diarrhœa, at the same age, there were 0'6 deaths, and consequently 99'4 recoveries, and so on for other ages.

TABLE XIII.—SUMMARY of Two Sets of SCOTCH RETURNS—No. I. from first appearance of CHOLERA in Scotland to 1st April 1854, and No. II. from 1st April 1854 to January 10th 1855.

Number of Parishes in Scotland.	Total Number of Parishes in which Cholera appeared.	Dates of first and last Cases in Returns No. I. & II.	Total Number of Cases.	Total Number of Deaths.
883	227*	{ 31st Aug. 1853 17th Nov. 1854 }	14,430*	6,848*

\* These numbers are larger than the combined totals in Returns No. I. and II. to the extent of 1 parish, 5 cases, and 4 deaths, reported to the Board of Supervision subsequent to the dispatch of Return No. I. to the General Board of Health on 18th May 1854.

*Board of Supervision,  
Edinburgh, 10th January 1855.*

W. S. WARREN,  
*Secretary.*

## SECOND SERIES.

TABLE I.—Showing the DEATHS in LONDON from CHOLERA and DIARRHŒA on each DAY from 1st July 1853 to 31st December 1854, both inclusive  
(C. denotes DEATHS from CHOLERA, and D. DEATHS from DIARRHŒA.)

Day of the Month.	1853.												1854.																							
	July.		Aug.		Sept.		Oct.		Nov.		Dec.		Jan.		Feb.		Mar.		April.		May.		June.		July.		Aug.		Sept.		Oct.		Nov.		Dec.	
	C.	D.	C.	D.	C.	D.	C.	D.	C.	D.	C.	D.	C.	D.	C.	D.	C.	D.	C.	D.	C.	D.	C.	D.	C.	D.	C.	D.	C.	D.	C.	D.	C.	D.		
1	—	6	—	20	1	21	11	14	15	10	—	10	—	6	7	3	—	6	—	5	7	—	4	—	—	6	72	31	389	41	70	13	9	—	2	
2	—	7	—	21	4	23	9	14	15	5	—	7	—	7	3	—	—	4	—	6	—	—	—	—	—	74	27	459	36	61	17	11	4	5		
3	—	8	—	23	5	24	10	13	12	7	—	8	—	1	—	—	—	2	—	1	—	—	—	—	—	78	28	329	49	66	13	4	—	6		
4	—	9	—	24	6	25	11	14	13	22	—	9	—	—	—	—	—	1	—	—	—	—	—	—	78	29	305	48	53	20	11	—	3			
5	—	7	—	18	3	31	7	13	10	3	—	1	—	—	—	—	—	—	—	—	—	—	—	—	101	26	267	47	60	16	4	7	—	8		
6	—	9	—	4	4	11	6	12	15	10	—	6	—	—	—	—	—	—	—	—	—	—	—	—	104	34	34	259	41	41	26	6	—	6		
7	—	12	—	7	1	16	7	9	13	7	—	8	—	—	—	—	—	—	—	—	—	—	—	—	78	28	335	35	37	15	2	8	—	3		
8	—	5	—	2	2	16	8	10	16	11	—	4	—	—	—	—	—	—	—	—	—	—	—	—	107	34	215	34	47	21	8	—	—	2		
9	—	10	—	22	7	16	10	8	12	2	—	4	—	—	—	—	—	—	—	—	—	—	—	—	96	25	259	39	37	24	6	5	—	5		
10	—	2	—	27	1	23	3	7	19	8	—	4	—	—	—	—	—	—	—	—	—	—	—	—	83	31	256	29	38	19	11	5	—	10		
11	—	9	—	22	3	6	7	8	15	9	—	4	—	—	—	—	—	—	—	—	—	—	—	—	105	49	233	52	29	11	3	6	—	3		
12	—	10	—	22	29	3	14	9	11	4	—	4	—	—	—	—	—	—	—	—	—	—	—	—	98	36	246	36	17	18	5	—	—	8		
13	—	11	—	5	25	8	10	7	12	10	—	4	—	—	—	—	—	—	—	—	—	—	—	—	108	33	263	43	24	18	7	—	—	2		
14	—	12	—	2	38	2	10	8	9	4	—	4	—	—	—	—	—	—	—	—	—	—	—	—	116	32	158	30	35	11	2	8	—	5		
15	—	14	—	21	4	15	3	8	12	9	—	4	—	—	—	—	—	—	—	—	—	—	—	—	111	31	203	30	32	17	1	5	—	4		
16	—	1	—	24	4	8	10	9	4	9	—	4	—	—	—	—	—	—	—	—	—	—	—	—	115	33	223	30	19	13	6	5	—	5		
17	—	16	—	19	8	11	14	11	10	3	—	4	—	—	—	—	—	—	—	—	—	—	—	—	125	36	190	39	26	14	1	1	—	4		
18	—	14	—	26	1	17	14	11	10	3	—	4	—	—	—	—	—	—	—	—	—	—	—	—	97	36	179	21	18	21	2	—	—	5		
19	—	18	—	3	27	5	11	7	9	4	—	4	—	—	—	—	—	—	—	—	—	—	—	—	118	38	167	23	25	18	4	—	—	3		
20	—	1	—	12	3	13	13	9	6	5	—	4	—	—	—	—	—	—	—	—	—	—	—	—	131	38	142	29	13	10	1	3	—	6		
21	—	10	—	5	31	7	16	9	5	3	—	4	—	—	—	—	—	—	—	—	—	—	—	—	131	39	159	26	13	11	2	3	—	5		
22	—	1	—	3	20	1	11	22	10	6	—	4	—	—	—	—	—	—	—	—	—	—	—	—	131	33	137	30	12	9	3	8	—	2		
23	—	16	—	8	13	11	14	7	6	11	—	4	—	—	—	—	—	—	—	—	—	—	—	—	140	41	129	28	12	9	—	5	—	4		
24	—	10	—	3	16	3	14	10	5	4	—	4	—	—	—	—	—	—	—	—	—	—	—	—	118	29	104	30	3	2	2	—	—	8		
25	—	17	—	3	14	6	9	6	9	4	—	4	—	—	—	—	—	—	—	—	—	—	—	—	115	36	106	26	4	9	—	5	—	4		
26	—	15	—	16	9	13	14	10	4	5	—	4	—	—	—	—	—	—	—	—	—	—	—	—	122	26	90	20	6	10	1	3	—	3		
27	—	22	—	3	16	11	14	12	4	8	—	4	—	—	—	—	—	—	—	—	—	—	—	—	141	41	88	25	5	4	7	1	—	5		
28	—	12	—	26	8	14	16	6	5	2	—	4	—	—	—	—	—	—	—	—	—	—	—	—	137	38	77	20	6	6	8	—	—	2		
29	—	19	—	2	35	6	12	14	8	3	—	4	—	—	—	—	—	—	—	—	—	—	—	—	187	29	66	17	3	9	—	—	—	1		
30	—	1	—	2	28	8	9	11	6	4	—	4	—	—	—	—	—	—	—	—	—	—	—	—	211	27	—	—	—	8	—	—	—	—		
31	—	18	—	25	1	—	25	8	4	5	—	4	—	—	—	—	—	—	—	—	—	—	—	—	44	20	40	—	9	—	4	—	—	—		
Total -	21	380	67	723	111	454	835	283	298	278	43	187	7	164	3	137	—	154	4	138	4	141	3	169	308	371	3,513	1,022	6,084	990	823	426	52	175	5	113

## SECOND SERIES—continued.

TABLE II.—DEATHS FROM CHOLERA and DIARRHŒA on each DAY of Five Months of the Year 1849.

Day of the Month.	June.			July.			August.			September.			October.		
	Chol.	Diarr.	Total.	Chol.	Diarr.	Total.	Chol.	Diarr.	Total.	Chol.	Diarr.	Total.	Chol.	Diarr.	Total.
1	1	1	2	19	12	31	142	29	171	250	42	292	41	16	57
2	2	7	9	37	14	51	123	36	159	282	36	320	55	20	75
3	2	5	7	30	10	40	98	29	127	316	37	353	31	13	44
4	—	2	2	20	3	23	138	27	165	336	42	378	22	20	42
5	3	2	5	21	10	31	127	30	157	314	41	355	28	16	44
6	4	3	7	29	8	37	96	25	121	297	44	341	20	16	36
7	10	3	13	35	14	49	110	29	139	284	34	318	23	19	42
8	7	9	16	34	12	46	128	38	166	311	41	352	13	11	24
9	7	5	12	45	10	55	104	20	124	299	34	333	11	11	22
10	5	12	17	67	11	78	124	32	156	264	47	311	16	10	26
11	9	8	17	58	12	70	137	23	180	226	39	265	14	9	23
12	5	7	12	46	14	60	166	30	196	217	46	238	7	12	19
13	9	3	12	79	16	95	214	21	235	217	38	255	9	6	15
14	5	7	12	91	15	106	204	32	236	156	43	199	5	4	9
15	7	5	12	97	24	121	185	30	215	153	46	199	7	7	14
16	10	7	17	121	24	145	173	35	208	120	41	161	3	10	13
17	7	1	8	79	28	107	151	25	176	131	33	164	3	7	10
18	12	1	13	84	28	112	155	42	187	109	30	139	2	9	11
19	6	7	13	101	27	128	168	41	209	119	30	154	2	7	9
20	8	2	10	120	31	151	214	40	254	89	25	114	1	7	8
21	8	7	15	126	29	155	193	37	230	69	26	95	4	6	10
22	6	4	10	109	34	143	158	27	185	75	30	105	1	5	6
23	8	3	11	104	45	149	180	51	231	71	29	100	2	7	9
24	14	10	24	95	35	130	203	32	235	62	29	91	3	4	7
25	10	1	11	123	32	155	240	27	267	63	24	87	—	7	7
26	16	3	19	114	39	153	220	46	266	46	18	64	4	8	12
27	23	8	31	109	33	142	249	29	278	57	18	75	2	6	8
28	24	4	28	117	28	145	280	28	308	38	20	58	5	7	12
29	27	6	33	121	24	145	238	39	277	41	20	61	2	4	6
30	24	7	31	172	24	196	243	35	278	44	27	71	1	6	7
31	—	—	—	152	29	181	187	28	215	—	—	—	—	5	5
Total	279	150	429	2,555	684	3,239	5,363	993	6,361	5,031	1,012	6,043	337	295	632



SECOND SERIES—*continued.*

TABLE III.—Showing the NUMBER of DEATHS by CHOLERA and DIARRHŒA in each DISTRICT of LONDON in the Year and half, 1853½ to 1854, distinguishing DEATHS in HOSPITALS.

No. of Districts.	Districts.	Deaths by		Deaths by		Total.
		Cholera, including Deaths in Hospitals.	Cholera in Hospitals.	Cholera, exclusive of Deaths in Hospitals.	Diarrhœa.	
	LONDON - - - - -	11,661	800	10,861	6,258	17,119
	West Districts :					
1	Kensington - - - - -	543	29	514	287	801
2	Chelsea - - - - -	309	—	309	153	467
3	St. George, Hanover Square - - -	303	64	239	130	369
4	Westminster - - - - -	443	63	380	207	587
5	St. Martin-in-the-Fields - - -	59	11	48	43	91
6	St. James, Westminster - - -	497	—	497	43	540
	North Districts :					
7	Marylebone - - - - -	397	123	269	349	613
8	Hampstead - - - - -	15	—	15	14	29
9	Pancras - - - - -	256	84	172	337	509
10	Islington - - - - -	116	—	116	249	365
11	Hackney - - - - -	91	—	91	127	218
	Central Districts :					
12	St. Giles - - - - -	115	—	115	141	256
13	Strand - - - - -	112	17	95	93	183
14	Holborn - - - - -	28	—	28	94	122
15	Clerkenwell - - - - -	67	—	67	127	194
16	St. Luke - - - - -	55	—	55	161	216
17	East London - - - - -	104	—	104	121	225
18	West London - - - - -	116	87	29	51	80
19	London, City - - - - -	79	—	79	60	139
	East Districts :					
20	Shoreditch - - - - -	266	—	266	298	564
21	Bethnal Green - - - - -	213	—	213	217	430
22	Whitechapel - - - - -	427	65	362	227	589
23	St. George-in-the-East - - -	175	—	175	156	331
24	Stepney - - - - -	436	—	436	323	759
25	Poplar - - - - -	218	—	218	137	355
	South Districts :					
26	St. Saviour, Southwark - - -	551	49	502	195	697
27	St. Olave, Southwark - - -	341	90	251	35	286
28	Bermondsey - - - - -	923	—	923	209	1,132
29	St. George, Southwark - - -	625	—	625	185	810
30	Newington - - - - -	741	—	741	196	937
31	Lambeth - - - - -	1,003	—	1,003	461	1,464
32	Wandsworth - - - - -	450	—	450	153	603
33	Camberwell - - - - -	531	—	531	180	761
34	Rotherhithe - - - - -	308	—	308	105	413
35	Greenwich - - - - -	613	113	500	308	803
36	Lewisham - - - - -	85	—	85	81	166

St. Bartholomew's Hospital, in the West London District, contained 540 inmates in 1851.  
 The portions of Guy's and St. Thomas's Hospitals, in the District of St. Olave, Southwark, contained 772 inmates in 1851.

SECOND SERIES—*continued.*

TABLE IV.—Showing the TOTAL NUMBER of DEATHS from CHOLERA and DIARRHOEA registered in each of the DISTRICTS and SUB-DISTRICTS in LONDON, from 1st July 1853 to 31st December 1854, both inclusive.

	District.	Registrar's Sub-District.	Cholera.	Diar.	Total.
<i>West Districts.</i>					
1	KENSINGTON - - -	Paddington, St. Mary - - -	33	47	80
		„ St. John - - -	71	52	123
		Kensington Town - - -	157	92	249
		Brompton - - -	48	29	77
		Hammersmith, St. Peter - - -	33	6	39
		„ St. Paul - - -	96	29	125
		Fulham - - -	105	32	137
2	CHELSEA - - -	Chelsea, South - - -	128	50	178
		„ North-west - - -	101	47	148
		„ North-east - - -	80	61	141
3	ST. GEORGE, HANOVER SQUARE.	Hanover Square - - -	21	26	47
		Mayfair - - -	32	13	45
		Belgrave - - -	250	91	341
4	WESTMINSTER - - -	St. John - - -	192	104	296
		St. Margaret - - -	251	103	354
5	ST. MARTIN IN FIELDS	Charing Cross - - -	41	16	57
		Long Acre - - -	18	27	45
6	ST. JAMES, WESTMINSTER.	Berwick Street - - -	200	17	217
		St. James Square - - -	20	6	26
		Golden Square - - -	277	20	297
<i>North Districts.</i>					
7	MARYLEBONE - - -	All Souls - - -	158	48	206
		Cavendish Square - - -	12	15	27
		Rectory - - -	99	65	164
		St. Mary - - -	44	46	90
		Christ Church - - -	49	98	147
		St. John - - -	35	77	112
8	HAMPSTEAD - - -	Hampstead - - -	15	14	29
9	PANCRAS - - -	Regent's Park - - -	25	51	76
		Tottenham Court - - -	90	64	154
		Gray's Inn Lane - - -	60	41	101
		Somers Town - - -	33	53	86
		Camden Town - - -	27	72	99
		Kentish Town - - -	21	56	77
10	ISLINGTON - - -	Islington West - - -	72	131	203
		„ East - - -	44	118	162
11	HACKNEY - - -	Stoke Newington - - -	5	6	11
		Stamford Hill - - -	9	14	23
		West Hackney - - -	26	34	60
		Hackney - - -	41	45	86
		South Hackney - - -	10	28	38
<i>Central Districts.</i>					
12	ST. GILES - - -	St. George, Bloomsbury - - -	13	32	45
		St. Giles, South - - -	63	62	125
		„ North - - -	39	47	86
13	STRAND - - -	St. Ann, Soho - - -	65	34	99
		St. Mary-le-Strand - - -	13	22	35
		St. Clement Danes - - -	34	37	71

SECOND SERIES—*continued.*TABLE IV.—Showing Total Number of Deaths from Cholera registered, &c.—*continued.*

	District.	Registrar's Sub-District.	Cholera.	Diar.	Total.
14	HOLBORN - - -	St. George Martyr - - St. Andrew, Holborn - Saffron Hill - - -	8 12 8	31 48 15	39 60 23
15	CLERKENWELL - -	St. James - - - Amwell - - - Pentonville - - - Goswell Street - - -	25 17 11 14	54 28 23 22	79 45 34 36
16	ST. LUKE - - -	Old Street - - - City Road - - - Whitecross Street - - Finsbury - - -	9 13 21 12	19 38 68 36	28 51 89 48
17	EAST LONDON - -	St. Botolph - - - Cripplelegate - - -	49 41	72 49	121 90
18	WEST LONDON - -	West London, North - "      South -	116 17	36 15	152 32
19	CITY OF LONDON -	London City, South West - "      North West - "      South - "      South East - "      North East -	15 8 19 24 10	9 9 10 11 21	24 17 29 35 31
<i>East Districts.</i>					
20	SHOREDITCH - - -	Holywell - - - St. Leonard - - - Hoxton, New Town - - "      Old Town - - Haggerstone, West - - "      East - - -	59 94 36 22 38 17	44 58 58 56 61 21	103 152 94 78 99 38
21	BETHNAL GREEN -	Hackney Road - - - Green - - - Church - - - Town - - -	57 81 32 43	46 79 43 49	103 160 75 92
22	WHITECHAPEL - -	Artillery - - - Spitalfields - - - Mile End, New Town - - Whitechapel, North - - "      Church - - Goodman's Fields - - Aldgate - - -	24 46 100 52 78 37 90	15 40 49 35 13 37 38	39 86 149 87 91 74 128
23	ST. GEORGE IN THE EAST.	St. Mary - - - St. Paul - - - St. John - - -	60 82 33	55 43 58	115 125 91
24	STEPNEY - - -	Shadwell - - - Ratcliff - - - Mile End, Old Town, Upper "      "      Lower Limehouse - - -	91 82 39 131 93	63 47 69 98 46	154 129 108 229 139
25	POPLAR - - -	Bow - - - Poplar - - -	81 137	59 78	140 215
			5,440	4,150	9,590

## SECOND SERIES--continued.

TABLE IV.—Showing Total Number of Deaths from Cholera registered, &amp;c.—continued.

	District.	Registrar's Sub-District.	Cholera.	Diar.	Total.
<i>South Districts.</i>					
26	ST. SAVIOUR, SOUTH-WARK.	Christ Church - - - St. Saviour - - -	127 424	48 147	175 571
27	ST. OLAVE, SOUTH-WARK.	St. Olave - - - - St. John - - - -	183 158	19 16	202 174
28	BERMONDSEY - -	St. James - - - St. Mary Magdalen - - Leather Market - -	388 275 260	64 73 72	452 348 332
29	ST. GEORGE, SOUTH-WARK.	Kent Road - - - Borough Road - - - London Road - - -	219 301 105	70 77 38	289 378 143
30	NEWINGTON - -	Trinity, Newington - - St. Peter, Walworth - - St. Mary - - -	224 419 98	69 111 16	293 530 114
31	LAMBETH - - -	Waterloo Road, 1st. - - 2nd. - - Lambeth Church, 1st. - - 2nd. - - Kennington, 1st. - - 2nd. - - Brixton - - - - Norwood - - - -	62 128 63 215 321 148 56 10	33 67 58 130 101 33 25 14	95 195 121 345 422 181 81 24
32	WANDSWORH - -	Clapham - - - - Battersea - - - - Wandsworth - - - - Putney - - - - Streatham - - - -	178 181 66 9 16	38 60 35 6 14	216 241 101 15 30
33	CAMBERWELL - -	Dulwich - - - - Camberwell - - - - Peckham - - - - St. George - - - -	- 256 187 138	1 71 58 50	1 327 245 188
34	ROTHERHITHE - -	Rotherhithe - - - -	308	105	413
35	GREENWICH - -	St. Paul, Deptford - - St. Nicholas, Deptford - - Greenwich, West - - East - - Woolwich Dockyard - - Arsenal - -	108 128 173 124 20 60	67 33 50 95 31 32	175 161 223 219 51 92
36	LEWISHAM - - -	Plumstead - - - - Eltham - - - - Lee - - - - Lewisham Village - - Sydenham - - - -	23 3 16 31 12	38 2 16 18 7	61 5 32 49 19
			6,221	2,108	8,329
DISTRICTS NORTH OF THE THAMES -			5,440	4,150	9,590
DISTRICTS SOUTH OF THE THAMES -			6,221	2,108	8,329
TOTAL DISTRICTS -			11,661	6,258	17,919



## SECOND SERIES--continued.

TABLE V.—SUMMARY TABLE, showing the NUMBER of DEATHS from CHOLERA and DIARRHOEA in the following HOSPITALS and WORKHOUSES of London, from the 1st day of July 1853 to the 31st of December 1854, both inclusive.

	District.	Sub-district.	Hospital, Workhouse, &c.	Cholera.	Diarrhoea.	Total.
<i>Districts North of the Thames:</i>						
1	KENSINGTON	Paddington, St. Mary	St. Mary's Hospital	29	-	29
			Paddington Workhouse	1	1	2
		Kensington Town	Kensington Workhouse	18	4	22
		Fulham	Earl's Court Workhouse*	4	10	14
			Fulham Workhouse	42	2	44
2	CHELSEA	Chelsea, North-west	Chelsea Workhouse	23	10	33
			St. George's Workhouse	18	8	26
3	ST. GEORGE, HANOVER SQUARE.	Mayfair	St. George, Hanover Square	19	4	23
		Belgrave	St. George's Hospital	64	5	69
4	WESTMINSTER	St. Margaret	Westminster Hospital	63	3	66
			Westminster Workhouse	49	20	69
5	ST. MARTIN-IN-THE-FIELDS.	Charing Cross	Charing Cross Hospital	11	-	11
			Charing Cross Workhouse	2	4	6
6	ST. JAMES, WESTMINSTER.	Golden Square	Workhouse, Golden Square	32	-	32
7	MARYLEBONE	All Souls	Middlesex Hospital	128	-	128
		Rectory	Marylebone Workhouse	46	14	60
8	HAMPSTEAD	Hampstead	Hampstead Workhouse	2	1	3
9	PANCRAS	Tottenham Court	University College Hospital	36	2	38
			Strand Workhouse	17	8	25
		Gray's Inn Lane	Royal Free Hospital	48	2	50
		Camden Town	Pancras Workhouse	15	5	20
10	ISLINGTON	Islington, West	Islington Workhouse	4	3	7
11	HACKNEY	Hackney	Hackney Workhouse	2	9	11
12	ST. GILES	St. Giles, South	St. Giles's Workhouse	35	5	40
13	STRAND	St. Clement Danes	King's College Hospital	17	1	18
14	HOLBORN	St. Andrew, Holborn	St. Andrew's Workhouse	-	9	9
15	CLERKENWELL	St. James	St. James's Workhouse	5	3	8
17	EAST LONDON	St. Botolph	St. Botolph Workhouse	10	18	28
18	WEST LONDON	West London, North	St. Bartholomew's Hospital	110	4	114
			Workhouse, Aldersgate	1	3	4
			Workhouse, West Street	-	4	4
20	SHOREDITCH	Hoxton New Town	St. Luke's Workhouse	17	14	31
		Haggerstone, West	Shoreditch Workhouse	13	18	31
21	BETHNAL GREEN	Green	Bethnal Green Workhouse	22	16	38
22	WHITECHAPEL	Mile End New Town	Workhouse	42	12	54
		Whitechapel, North	Workhouse	29	6	35
		Whitechapel Church	London Hospital	65	4	69
23	ST. GEORGE IN THE EAST.	St. John	Workhouse	2	10	12
24	STEPNEY	Shadwell	Shadwell Workhouse	3	9	12
		Ratcliff	Ratcliff Workhouse	-	4	4
		Mile End Old Town,	Mile End Workhouse	70	5	75
		Lower, &c.	London City Workhouse	-	3	3
		Limehouse	Limehouse Workhouse	-	-	-
25	POPLAR	Bow	London City Workhouse	-	4	4
		Poplar	Poplar Workhouse	26	13	39
<i>Districts South of the Thames:</i>				1,140	280	1,420
26	ST. SAVIOUR, SOUTHWARK.	Christ Church	St. Saviour's Workhouse	16	7	23
		St. Saviour	Guy's Hospital	45	-	45
			St. Thomas's Hospital	4	2	6
27	ST. OLAVE, SOUTHWARK.	St. Olave	Guy's Hospital	31	6	37
			St. Thomas's Hospital	59	10	69
		St. John	St. John's Workhouse	16	4	20
28	BERMONDSEY	St. Mary Magdalen	Bermondsey Workhouse	38	9	47
29	ST. GEORGE, SOUTHWARK.	Borough Road	Workhouse, Mint Street	80	20	100
30	NEWINGTON	St. Peter, Walworth	Workhouse	22	4	26
31	LAMBETH	Lambeth Church, 2d	Workhouse	59	43	102
		Norwood	Workhouse	1	3	4
32	WANDSWORTH	Battersea	Workhouse	12	5	17
33	CAMBERWELL	Camberwell	Camberwell Workhouse	68	5	73
34	ROTHERHITHE	Rotherhithe	Rotherhithe Workhouse	4	9	13
35	GREENWICH	Greenwich, West	Dreadnought Hospital	113	3	116
		Do, East	Workhouse	29	32	61
36	LEWISHAM	Lewisham Village	Workhouse	10	-	10
				607	162	769
DISTRICTS NORTH OF THE THAMES				1,140	280	1,420
DISTRICTS SOUTH OF THE THAMES				607	162	769
TOTAL DISTRICTS				1,747	442	2,189

\* This Workhouse belongs to the District of Westminster.

## SECOND SERIES—continued.

TABLE VI.—MORTALITY from CHOLERA and DIARRHŒA, 1853½-54, in the SUB-DISTRICTS of LONDON. (Certain Corrections are introduced for DEATHS in HOSPITALS and WORKHOUSES.)

No. of Sub-district.	District and Sub-district.	Deaths from Cholera and Diarrhœa to 10,000 persons living.		No. of Sub-district.	District and Sub-district.	Deaths from Cholera and Diarrhœa to 10,000 persons living.	
		Cholera.	Diarrhœa.			Cholera.	Diarrhœa.
	1. KENSINGTON.				10. ISLINGTON.		
1	St. Mary, Paddington - -	2	20	1	Islington, West - - -	13	22
2	St. John, Paddington - -	26	15	2	Islington, East - - -	9	21
3	Kensington Town - - -	49	27		11. HACKNEY.		
4	Brompton - - - -	34	17				
5	St. Peter, Hammersmith -	83	12	1	Stoke Newington - - -	11	12
6	St. Paul, Hammersmith -	80	20	2	Stamford Hill - - - -	17	25
7	Fulham - - - - -	59	25	3	West Hackney - - - -	13	15
	2. CHELSEA.			4	Hackney - - - - -	19	19
				5	South Hackney - - - -	12	31
1	Chelsea, South - - -	76	24		12. ST. GILES.		
2	Chelsea, North-west - -	38	24	1	St. George, Bloomsbury -	12	19
3	Chelsea, North-east - -	43	27	2	St. Giles, South - - -	22	31
	3. ST. GEORGE, HANOVER SQUARE.			3	St. Giles, North - - -	34	27
1	Hanover Square - - - -	12	13		13. STRAND.		
2	May Fair - - - - -	12	10	1	St. Anne, Soho - - - -	40	19
3	Belgrave - - - - -	49	21	2	St. Mary-le-Strand - - -	12	19
	4. WESTMINSTER.			3	St. Clement Danes - - -	12	24
1	St. John - - - - -	63	28		14. HOLBORN.		
2	St. Margaret - - - -	54	33	1	St. George the Martyr - -	4	16
	5. ST. MARTIN-IN-THE-FIELDS.			2	St. Andrew, Eastern - - -	9	33
				3	Saffron Hill - - - - -	6	11
1	Charing Cross - - - -	26	13		15. CLERKENWELL.		
2	Long Acre - - - - -	16	22	1	St. James - - - - -	12	24
	6. ST. JAMES, WESTMINSTER.			2	Anwell - - - - -	11	17
				3	Pentonville - - - - -	9	18
1	Berwick Street - - -	211	16	4	Goswell Street - - - -	9	14
2	St. James's Square - -	20	5		16. ST. LUKE.		
3	Golden Square - - -	197	14	1	Old Street - - - - -	9	17
	7. MARYLEBONE.			2	City Road - - - - -	8	22
				3	Whitecross Street - - -	16	49
1	All Souls - - - - -	13	16	4	Finsbury - - - - -	10	27
2	Cavendish Square - - -	10	10		17. EAST LONDON (CITY).		
3	Rectory - - - - -	25	23				
4	St. Mary - - - - -	25	20	1	St. Botolph - - - - -	19	29
5	Christchurch - - - -	18	27	2	Cripplegate - - - - -	23	23
6	St. John - - - - -	13	22		18. WEST LONDON (CITY).		
	8. HAMPSTEAD.						
1	Hampstead - - - - -	13	11	1	West London, North - -	5	29
	9. PANCRAS.			2	West London, South - -	12	9
					19. LONDON CITY.		
1	Regents Park - - - -	10	15	1	London City, South-west -	18	10
2	Tottenham Court - - -	17	22	2	London City, North-west -	7	8
3	Grays Inn Lane - - -	6	15	3	London City, South - - -	18	9
4	Somers Town - - - -	11	14	4	London City, South-east -	24	10
5	Camden Town - - - -	7	30	5	London City, North-east -	8	16
6	Kentish Town - - - -	9	18				

SECOND SERIES—*continued.*TABLE VI.—Mortality from Cholera and Diarrhœa, 1853-4, in the Sub-Districts of London, &c.—*continued.*

No. of Sub-district.	District and Sub-district.	Deaths from Cholera and Diarrhoea to 10,000 persons living.		No. of Sub-district.	District and Sub-district.	Deaths from Cholera and Diarrhoea to 10,000 persons living.	
		Cholera.	Diarrhoea.			Cholera.	Diarrhoea.
20. SHOREDITCH.							
1	Holywell - - - - -	41	25	1	29. ST. GEORGE, SOUTH-WARK.		
2	St. Leonard - - - - -	57	29				
3	Hoxton New Town - - -	8	22				
4	Hoxton Old Town - - -	14	30				
5	Haggerstone, West - - -	12	25				
6	Haggerstone, East - - -	15	16				
21. BETHNAL GREEN.							
1	Hackney Road - - - - -	27	18	1	Trinity, Newington - -	115	32
2	Green - - - - -	27	30	2	St. Peter, Walworth - -	136	34
3	Church - - - - -	16	18	3	St. Mary - - - - -	74	11
4	Town - - - - -	24	23	30. NEWINGTON.			
22. WHITECHAPEL.							
1	Artillery - - - - -	46	22	1	Waterloo Road, 1st - -	49	23
2	Spitalfields - - - - -	40	26	2	Waterloo Road, 2d - -	78	36
3	Mile End New Town - - -	50	32	3	Lambeth Church, 1st - -	39	32
4	Whitechapel, North - - -	25	28	4	Lambeth Church, 2d - -	63	46
5	Whitechapel Church - - -	21	16	5	Kennington, 1st - - -	135	37
6	Goodmans Fields - - - -	38	29	6	Kennington, 2d - - -	81	16
7	Aldgate - - - - -	107	34	7	Brixton - - - - -	39	15
23. ST. GEORGE-IN-THE-EAST.							
1	St. Mary - - - - -	34	29	8	Norwood - - - - -	27	32
2	St. Paul - - - - -	41	20	31. LAMBETH.			
3	St. John - - - - -	33	53	1	Waterloo Road, 1st - -	49	23
24. STEPNEY.							
1	Shadwell - - - - -	69	37	2	Waterloo Road, 2d - -	78	36
2	Ratcliff - - - - -	64	28	3	Lambeth Church, 1st - -	39	32
3	Mile End Old Town, Upper -	16	22	4	Lambeth Church, 2d - -	63	46
4	Mile End Old Town, Lower -	26	33	5	Kennington, 1st - - -	135	37
5	Linehouse - - - - -	50	19	6	Kennington, 2d - - -	81	16
25. POPLAR.							
1	Bow - - - - -	44	26	7	Brixton - - - - -	39	15
2	Poplar - - - - -	42	25	8	Norwood - - - - -	27	32
26. ST. SAVIOUR, SOUTH-WARK.							
1	Christchurch - - - - -	75	29	32. WANDSWORTH.			
2	St. Saviour - - - - -	208	73	1	Clapham - - - - -	109	21
27. ST. OLAVE, SOUTHWARK.							
1	St. Olave - - - - -	163	26	2	Battersea - - - - -	152	49
2	St. John, Horsleydown - -	140	14	3	Wandsworth - - - - -	70	34
28. BERMONDSEY.							
1	St. James - - - - -	201	29	4	Putney - - - - -	18	11
2	St. Mary Magdalen - - -	169	47	5	Streatham - - - - -	19	15
3	Leather Market - - - -	179	44	33. CAMBERWELL.			
29. ST. GEORGE, SOUTH-WARK.							
1	Holywell - - - - -	41	25	1	Dulwich - - - - -	—	6
2	St. Leonard - - - - -	57	29	2	Camberwell - - - - -	120	37
3	Hoxton New Town - - -	8	22	3	Peckham - - - - -	101	26
4	Hoxton Old Town - - -	14	30	4	St. George - - - - -	94	28
5	Haggerstone, West - - -	12	25	34. ROTHERHITHE.			
6	Haggerstone, East - - -	15	16	1	Rotherhithe - - - - -	171	54
30. NEWINGTON.							
1	Trinity, Newington - -	115	32	35. GREENWICH.			
2	St. Peter, Walworth - -	136	34	1	St. Paul, Deptford - -	45	24
3	St. Mary - - - - -	74	11	2	St. Nicholas, Deptford -	206	46
31. LAMBETH.							
1	Waterloo Road, 1st - -	49	23	3	Greenwich, West - - -	35	26
2	Waterloo Road, 2d - -	78	36	4	Greenwich, East - - -	62	54
3	Lambeth Church, 1st - -	39	32	5	Woolwich Dockyard - -	12	16
4	Lambeth Church, 2d - -	63	46	6	Woolwich Arsenal - - -	43	20
5	Kennington, 1st - - -	135	37	36. LEWISHAM.			
6	Kennington, 2d - - -	81	16	1	Plumstead - - - - -	16	22
7	Brixton - - - - -	39	15	2	Eltham - - - - -	12	8
8	Norwood - - - - -	27	32	3	Lee - - - - -	21	18
32. WANDSWORTH.							
1	Clapham - - - - -	109	21	4	Lewisham Village - - -	41	28
2	Battersea - - - - -	152	49	5	Sydenham - - - - -	29	13
3	Wandsworth - - - - -	70	34	37. GREENWICH.			
4	Putney - - - - -	18	11	1	St. Paul, Deptford - -	45	24
5	Streatham - - - - -	19	15	2	St. Nicholas, Deptford -	206	46
33. CAMBERWELL.							
1	Dulwich - - - - -	—	6	3	Greenwich, West - - -	35	26
2	Camberwell - - - - -	120	37	4	Greenwich, East - - -	62	54
3	Peckham - - - - -	101	26	5	Woolwich Dockyard - -	12	16
4	St. George - - - - -	94	28	6	Woolwich Arsenal - - -	43	20
34. ROTHERHITHE.							
1	Rotherhithe - - - - -	171	54	36. LEWISHAM.			
35. GREENWICH.							
1	St. Paul, Deptford - -	45	24	1	Plumstead - - - - -	16	22
2	St. Nicholas, Deptford -	206	46	2	Eltham - - - - -	12	8
3	Greenwich, West - - -	35	26	3	Lee - - - - -	21	18
4	Greenwich, East - - -	62	54	4	Lewisham Village - - -	41	28
5	Woolwich Dockyard - -	12	16	5	Sydenham - - - - -	29	13
6	Woolwich Arsenal - - -	43	20	37. GREENWICH.			
36. LEWISHAM.							
1	Plumstead - - - - -	16	22	1	St. Paul, Deptford - -	45	24
2	Eltham - - - - -	12	8	2	St. Nicholas, Deptford -	206	46
3	Lee - - - - -	21	18	3	Greenwich, West - - -	35	26
4	Lewisham Village - - -	41	28	4	Greenwich, East - - -	62	54
5	Sydenham - - - - -	29	13	5	Woolwich Dockyard - -	12	16

The 800 deaths from cholera in hospitals have been distributed over all the sub-districts, and where many deaths from cholera occurred in workhouses, a proportion has been distributed over the other sub-districts of the union in which the workhouse is situated; the population used for this calculation has been corrected to the middle (1st July) of 1854.

SECOND SERIES—*continued.*

TABLE VII.—Showing the AGES of those who DIED from DIARRHŒA in the NORTH and SOUTH DISTRICTS of the METROPOLIS, distinguishing Males and Females, from 1st July 1853 to 31st December 1854.

Age.	North Districts.			South Districts.			North and South Districts.			Age.
	Male.	Female.	Total.	Male.	Female.	Total.	Male.	Female.	Total.	
0—	1,071	900	1,971	427	356	783	1,498	1,256	2,754	0—
1—	303	303	606	181	175	356	484	478	962	1—
2—	113	84	197	73	59	132	186	143	329	2—
3—	38	34	72	27	26	53	65	69	125	3—
4—	29	26	55	13	12	25	42	38	80	4—
5—	56	65	121	37	22	59	93	87	180	5—
10—	20	19	39	12	10	22	32	29	61	10—
15—	18	27	45	11	11	22	29	38	67	15—
20—	18	29	47	7	22	29	25	51	76	20—
25—	19	37	56	11	23	34	30	60	90	25—
30—	24	30	54	24	21	45	48	51	99	30—
35—	23	37	60	16	23	39	39	60	99	35—
40—	28	32	60	7	20	27	35	52	87	40—
45—	26	22	48	11	27	38	37	49	86	45—
50—	43	40	83	17	22	39	60	62	122	50—
55—	25	47	72	20	23	43	45	70	115	55—
60—	49	61	110	41	36	77	90	97	187	60—
65—	52	60	112	30	34	64	82	94	176	65—
70—	58	72	130	39	48	87	97	120	217	70—
75—	42	76	118	34	29	63	76	105	181	75—
80—	20	36	56	16	26	42	36	62	98	80—
85—	8	21	29	7	12	19	15	33	48	85—
90—	—	5	5	3	4	7	3	9	12	90—
95—	1	1	2	—	3	3	1	4	5	95—
?	—	2	2	—	—	—	—	2	2	?
Total -	2,084	2,066	4,150	1,064	1,044	2,108	3,148	3,110	6,258	

TABLE VIII.—Showing the AGES of those who DIED in LONDON from CHOLERA in the DISTRICTS NORTH and SOUTH of the THAMES, separately and collectively, between the 1st July 1853 and the 31st December 1854.

Age.	Districts North of the Thames.			Districts South of the Thames.			North and South Districts.			Age.
	Male.	Female.	Total.	Male.	Female.	Total.	Male.	Female.	Total.	
0—	103	104	207	87	53	140	190	157	347	0—
1—	106	85	191	159	162	321	265	247	512	1—
2—	107	102	209	159	143	302	266	245	511	2—
3—	101	80	181	126	115	241	227	195	422	3—
4—	87	53	140	115	90	205	202	143	345	4—
5—	265	223	488	292	271	563	557	494	1,051	5—
10—	136	98	234	114	104	218	250	202	452	10—
15—	95	92	187	112	123	235	207	215	422	15—
20—	160	142	302	159	203	362	319	345	664	20—
25—	180	162	342	200	223	423	380	385	765	25—
30—	194	215	409	195	203	461	389	481	870	30—
35—	214	219	433	225	276	501	439	495	934	35—
40—	199	241	440	178	239	417	377	480	857	40—
45—	183	176	359	176	188	364	359	364	723	45—
50—	160	192	352	133	167	300	293	359	652	50—
55—	129	131	260	123	169	292	252	300	552	55—
60—	104	167	271	132	177	309	236	344	580	60—
65—	81	108	189	89	129	218	170	237	407	65—
70—	50	78	128	52	99	151	102	177	279	70—
75—	22	47	69	45	74	119	67	121	188	75—
80—	13	19	32	18	29	47	31	48	79	80—
85—	1	8	9	4	15	19	5	23	28	85—
90—	—	1	1	—	3	3	—	4	4	90—
95—	—	2	2	—	—	—	—	2	2	95—
?	1	4	5	10	—	10	11	4	15	?
Total -	2,691	2,749	5,440	2,903	3,318	6,221	5,594	6,067	11,661	



## SECOND SERIES—continued.

TABLE IX.—Showing the DEATHS from CHOLERA and DIARRHŒA in LONDON at Twelve different AGES, in the Years 1849 and 1853½–54; also the Enumerated Population of 1841 and 1851.

Ages.	Deaths by Cholera.				Deaths by Diarrhœa.				Population.	
	1849.	1853½–54.	Total.	Mean.	Mean.	Total.	1849.	1853½–54.	1841.	1851.
All Ages	14,137	11,661	25,798	12,899		10,157	3,899	6,258	1,948,417	2,362,236
0—	1,969	2,142	4,111	2,056		6,765	2,515	4,251	231,018	293,562
5—	1,298	1,056	2,354	1,177		300	120	181	194,199	243,648
10—	679	452	1,131	565		100	39	61	178,023	216,369
15—	1,334	1,086	2,420	1,210		208	65	143	393,528	455,095
25—	2,036	1,637	3,673	1,837		277	88	189	366,416	428,123
35—	2,108	1,793	3,901	1,950		328	142	186	256,886	308,949
45—	1,861	1,376	3,237	1,619		343	135	208	166,641	208,363
55—	1,475	1,132	2,607	1,303		525	222	302	94,830	122,946
65—	974	686	1,660	830		705	311	393	48,939	62,608
75—	368	267	635	318		493	214	279	15,574	19,845
85—	35	32	67	33		106	46	60	2,218	2,578
95 & upwards	—	2	2	1			2	5	145	150

SECOND SERIES—*continued.*

TABLE X.—Showing the MEAN RATE of MORTALITY at Twelve different AGES from CHOLERA and DIARRHŒA in London in the Two Epidemics of 1849 and 1853–54.

Age.	Mean of the Deaths, in each Epidemic, to 10,000 Persons, living at each Age from	
	CHOLERA.	DIARRHŒA.
	Mean of 1849 and 1853–54.	Mean of 1849 and 1853–54.
All Ages	53·7	21·1
0—	68·8	113·3
5—	47·5	6·1
10—	25·7	2·3
15—	26·1	2·2
25—	42·2	3·2
35—	62·1	5·2
45—	76·4	8·1
55—	104·2	21·0
65—	130·3	55·3
75—	157·3	122·1
85—	127·7	202·1
95 & upwards	65·5	229·4

The Table may be read thus:—To 10,000 Persons living at the Age 25 and under 35, 42 died of Cholera and 3 in 10,000 of Diarrhœa, on an average in each epidemic; and the mortality in the *two* epidemics was at the rate of 84·4 by Cholera, and 6·4 by Diarrhœa in 10,000 of the population living at that age.

## THIRD SERIES.

TABLE I.—MORTALITY from CHOLERA in the 36 Districts of LONDON during the Two Epidemics of 1849 and 1853½–1854; with the Elevation, Density, and other Conditions of the Population.

Company by which Water is chiefly supplied.	DISTRICTS.	Annual Value of Houses.	Mean Elevation in Feet (above Trinity High-water Mark).*	Area in Acres.	Popu- lation, 1851.	Density (1851).	Mortality from Cholera to every 10,000 Inhabitants.		Mortality from Diarrhoea to every 10,000 Inhabitants.	
						Persons to an Acre.	In the Year 1849.	In the Years 1853½ to end of 1854.	In the Year 1849.	In the Years 1853½ to end of 1854.
	LONDON - - - -	£ 40	39	78,029	2,362,236	30	62	46	17	25
J. W.	Kensington HL - - -	43	40	7,374 <i>w</i>	120,004	16	24	38	13	20
C.	Chelsea WH - - -	29	12	865 <i>w</i>	56,538	65	46	51	17	25
J. C.	St. George, Hanover Sq. H	110	34	1,161 <i>w</i>	73,230	63	18	33	10	17
C.	Westminster HHPP -	36	3	917 <i>w</i>	65,609	72	68	57	17	30
N.	St. Martin in the Fields. H	119	38	305 <i>w</i>	24,640	81	37	20	11	17
J. N.	St. James, Westminster -	128	58	164	36,406	222	16	142	11	12
W.	Marylebone H - - -	71	87	1,509	157,696	105	17	17	15	21
N. W.	Hampstead - - -	40	350	2,252	11,986	5	8	12	10	11
J. H. W.	Pancras WHH - - -	41	73	2,716	166,956	61	22	10	14	19
N.	Islington HHPP - - -	35	94	3,127	95,329	30	22	11	13	22
N. E.	Hackney H - - -	25	53	3,929	58,429	15	25	15	17	20
N.	St. Giles - - -	60	68	245	54,214	221	53	22	14	26
N.	Strand H - - -	66	50	174 <i>w</i>	44,460	256	35	22	14	21
N.	Holborn - - -	52	53	196	46,621	238	35	6	12	20
N.	Clerkenwell PP - - -	33	65	380	64,778	170	19	10	14	19
N.	St. Luke L - - -	28	51	220	54,055	246	34	10	18	29
N. E.	East London P - - -	38	40	153	44,406	290	45	+23	13	26
N.	West London WHPPP†	65	29	136 <i>w</i>	28,790	212	96	+10	22	18
N.	City of London - - -	117	31	434 <i>w</i>	55,932	129	38	+14	8	11
N. E.	Shoreditch WL - - -	20	48	646	109,257	169	76	23	18	25
E.	Bethnal Green L - - -	9	38	760	90,193	119	90	23	24	23
E.	Whitechapel H - - -	26	32	406 <i>w</i>	79,759	196	64	45	20	27
E.	St. George in the East -	32	21	243	48,376	199	42	36	15	31
E.	Stepney H - - -	20	21	1,257 <i>w</i>	110,775	88	47	38	17	27
E.	Poplar WL - - -	44	8	2,918 <i>w</i>	47,162	16	71	42	17	25
S. L.	St. Saviour, Southwk. HH	36	3	250 <i>w</i>	35,731	143	153	142	32	53
S. L.	St. Olave, Southwk. HH	35	4	169 <i>w</i>	19,375	115	181	140	25	19
S.	Bermondsey - - -	18	0	688 <i>w</i>	48,128	70	161	179	33	39
S. L.	St. George, Southwk. LLP	22	0	282	51,824	184	164	121	25	34
L. S.	Newington P - - -	22	-1	624	64,816	104	144	112	21	29
L. S.	Lambeth - - -	28	3	4,015 <i>w</i>	139,325	35	120	70	20	31
S.	Wandsworth L - - -	29	24	11,695 <i>w</i>	50,764	4	100†	85	15	28
S. L.	Camberwell LL - - -	25	4	4,342	54,667	13	97	99	17	30
S. K.	Rotherhithe - - -	23	0	886 <i>w</i>	17,805	21	205	165	26	54
K.	Greenwich HHHP - - -	22	8	5,367 <i>w</i>	99,365	19	75	49	21	29
K. L.	Lewisham W - - -	27	28	17,224 <i>w</i>	34,835	2	30	22	17	20

\* The Elevation of the Districts has been deduced for these new Tables by multiplying the "numbers of the population" returned at the Census of 1851 into the "elevation" of each Sub-District, and dividing the sum of these by the population of the District. Thus the mean elevation of the ground on which the population resides is found with tolerable accuracy. The elevation for the Districts of Lambeth, Greenwich, and Lewisham is taken from a Return furnished by Colonel Dawson.

† The three districts of the City of London, the East London, and the West London, including St. Bartholomew's Hospital, comprise the City of London, within and without the walls. Mr. Simon, the health officer of the City, has ascertained that the deaths from cholera properly belonging to these districts were 213, 197, 278 in 1849, and 79, 104, and 29 in 1854; his numbers have been adopted in the calculations. A similar correction is required of the mortality of St. Olave's district, where many persons from other districts died of cholera in St. Thomas's Hospital.

‡ Excluding the deaths which occurred in Drouet's Asylum for Infant Paupers, the mortality of the Wandsworth District was at the rate of 72 deaths to 10,000 persons living.

\*.\* The areas marked thus (*w*) include portions of the River Thames.

The several Water Companies are designated by letters:—thus, the New River Company by N; the Grand Junction by J; Chelsea by C; West Middlesex by W; East London by E; Hampstead by H; Southwark by S; Lambeth by L; and Kent by K.

THIRD SERIES—*continued.*

TABLE II.—MEAN ELEVATION and MEAN MORTALITY by CHOLERA and DIARRHŒA in LONDON DISTRICTS of different Degrees of DENSITY.

Density. Persons to an Acre.	Mean Elevation in Feet.	Mean Mortality to 10,000 Persons.	
		Cholera.	Diarrhœa.
1 — 10	66	37	21
10 — 30	30	50	26
30 — 50	15	71	23
50 — 100	33	53	25
100 — 150	22	64	28
150 — 200	40	42	21
200 — 250	42	34	23
250 — 300	44	27	25
300 — 400	57	28	24
400 and ups.*	56	80	33

The Table may be read thus : — In districts containing 50 and under 100 persons to an acre, the mean elevation is 33 feet, and the mean mortality by Cholera was 53 in 10,000; by Diarrhœa 25 in 10,000 in London in the one and a half years 1853½–1854.

\* Including the sub-districts of Whitecross Street, St. Andrew Eastern, and Berwick Street, St. James's, to the latter of which exclusively the high mortality is due.



## THIRD SERIES—continued.

TABLE III.—Of the SUB-DISTRICTS of LONDON arranged according to the DENSITY of the POPULATION, with the MEAN ELEVATION above TRINITY HIGH-WATER MARK, and MORTALITY from CHOLERA in the  $1\frac{1}{2}$  years 1853 $\frac{1}{2}$ –1854.

SUB-DISTRICTS.	DISTRICTS.	Density. Persons to an Acre.	Elevation in Feet.	Mortality by Cholera and Diarrhoea in the $1\frac{1}{2}$ years 1853 $\frac{1}{2}$ –1854 to 10,000 persons living.	
				Cholera.	Diarrhoea.
Dulwich L - - -	Camberwell - - -	1	68	-	6
Eltham - - -	Lewisham - - -	1	?	12	8
Putney - - -	Wandsworth - - -	2	12	18	11
Sydenham - - -	Lewisham - - -	2	188 $\frac{1}{2}$	29	13
Lewisham Village Ww	Lewisham - - -	2	16	41	23
Plumstead - - -	Lewisham - - -	3	?	16	22
Streatham L - - -	Wandsworth - - -	3	72	19	15
Lee - - -	Lewisham - - -	4	42	21	18
Norwood WL - - -	Lambeth - - -	4	128 $\frac{1}{2}$	27	32
Wandsworth P - - -	Wandsworth - - -	4	12	70	34
Hampstead W - - -	Hampstead - - -	5	350	13	11
Battersea W - - -	Wandsworth - - -	5	3	152	49
Hammersmith, St. Paul - - -	Kensington - - -	6	8	80	20
Fulham w L - - -	Kensington - - -	6	6	59	25
Stoke Newington - - -	Hackney - - -	8	72	11	12
Stamford Hill - - -	Hackney - - -	9	76	17	25
Brixton - - -	Lambeth - - -	10	56	39	15
Greenwich, East W - - -	Greenwich - - -	10	7	62	54
Hackney W - - -	Hackney - - -	12	44	19	19
Clapham - - -	Wandsworth - - -	13	21	109	21
Bow w - - -	Poplar - - -	13	15	44	26
Camberwell W - - -	Camberwell - - -	13	5	120	37
Kentish Town - - -	Pancras - - -	14	110	9	18
Woolwich Arsenal P - - -	Greenwich - - -	14	?	43	20
St. Paul, Deptford - - -	Greenwich - - -	15	10	45	24
Peckham - - -	Camberwell - - -	17	4	101	26
South Hackney - - -	Hackney - - -	18	44	12	31
Poplar W - - -	Poplar - - -	19	3	42	25
Rotherhithe W - - -	Rotherhithe - - -	20	0	171	54
Paddington, St. Mary W - - -	Kensington - - -	21	82	2	20
Brompton H - - -	Kensington - - -	21	12	34	17
Hammersmith, St. Peter - - -	Kensington - - -	22	4	83	12
Kensington Town WL - - -	Kensington - - -	23	28	49	27
Islington East - - -	Islington - - -	25	88	9	21
LONDON - - -	LONDON - - -	30	39	46	25
Woolwich Dockyard - - -	Greenwich - - -	35	?	12	16
Kennington, 2d part - - -	Lambeth - - -	37	8	81	16
St. George - - -	Camberwell - - -	37	-3	94	28
Islington West WHP - - -	Islington - - -	39	100	13	22
West Hackney - - -	Hackney - - -	40	55	13	15
St. James - - -	Bermondsey - - -	42	-1	201	29
Hanover Square - - -	St. George, Hanover Square - - -	45	64	12	13
St. Nicholas, Deptford - - -	Greenwich - - -	47	4	206	46
Charing Cross WH - - -	St. Martin-in-the-Fields - - -	48	17	26	13
St. Margaret WHP - - -	Westminster - - -	48	4	54	33
Chelsea, South - - -	Chelsea - - -	52	10	76	24
Kennington, 1st part - - -	Lambeth - - -	53	4	135	37
St. John - - -	Marylebone - - -	55	124	13	22
Mile End Old Town, Lower W - - -	Stepney - - -	55	23	26	33
Greenwich, West P - - -	Greenwich - - -	58	12	35	26
Green W - - -	Bethnal Green - - -	60	36	27	30
Paddington, St. John H - - -	Kensington - - -	65	76	26	15
Christchurch - - -	Marylebone - - -	65	92	18	27
Belgrave H - - -	St. George, Hanover Square - - -	69	12	40	21
Chelsea, North-cast L - - -	Chelsea - - -	70	13	43	27
Regent's Park - - -	Pancras - - -	75	87	10	15
Chelsea, North-west Ww - - -	Chelsea - - -	83	12	38	24
Limchouse W - - -	Stepney - - -	86	10	50	19
St. Mary - - -	Newington - - -	87	-1	74	11
Pentonville - - -	Clerkenwell - - -	89	84	9	18
Lambeth Church, 1st part - - -	Lambeth - - -	89	2	39	32
Shadwell W - - -	Stepney - - -	90	7	69	37
Haggerstone, East - - -	Shoreditch - - -	91	52	15	16
St. Peter, Walworth W - - -	Newington - - -	93	-2	136	34
May Fair W - - -	St. George, Hanover Square - - -	95	56	12	10
St. Mary Magdalen W - - -	Bermondsey - - -	98	0	169	47
Aldgate - - -	Whitechapel - - -	99	19	107	34

## THIRD SERIES—continued.

TABLE III.—Of the Sub-Districts of London, &amp;c.—continued.

SUB-DISTRICTS.	DISTRICTS.	Density. Persons to an Acre.	Elevation in Feet.	Mortality by Cholera and Diarrhoea in the 1½ Years 1853½-1854 to 10,000 persons living.	
				Cholera.	Diarrhoea.
City of London, South-east	City of London	103	21	24	10
St. John W	St. George-in-the-East	103	2	33	53
St. Olave H	St. Olave, Southwark	107	6	163	26
City of London, South	City of London	115	21	18	9
Ratcliff W	Stepney	115	18	64	28
St. John, Horsleydown W	St. Olave, Southwark	121	2	140	14
Camden Town W	Paneras	123	62	7	30
St. Saviour H	St. Saviour, Southwark	127	4	208	73
Waterloo Road, 2d part	Lambeth	129	2	78	36
Cavendish Square	Marylebone	130	73	10	10
St. John HP	Westminster	132	2	63	28
City of London, South-west	City of London	137	21	18	10
St. George, Bloomsbury	St. Giles	138	71	12	19
City of London, North-east	City of London	139	44	8	16
Lambeth Church, 2d part W	Lambeth	144	1	63	46
Trinity P	Newington	147	-1	115	32
Hoxton Old Town	Shoreditch	150	52	14	30
Haggerstone, West W	Shoreditch	154	52	12	25
Mile End Old Town, Upper	Stepney	155	32	16	22
Waterloo Road, 1st part	Lambeth	155	3	49	23
London Road	St. George, Southwark	159	0	71	21
City of London, North-west	City of London	165	44	7	8
Church	Bethnal Green	165	36	16	18
Golden Square W	St. James, Westminster	166	68	197	14
Leather Market	Bermondsey	166	0	179	44
Whitechapel Church H	Whitechapel	163	32	21	16
Christchurch W	St. Saviour, Southwark	169	2	75	29
Hackney Road	Bethnal Green	170	44	27	18
Gray's Inn Lane II	Paneras	171	52	6	15
St. Mary-le-Strand	Strand	173	48	12	19
Kent Road	St. George, Southwark	173	-1	142	37
Goswell Street	Clerkenwell	174	78	9	14
West London, South	West London	178	24	12	9
Hoxton New Town W	Shoreditch	181	52	8	22
St. George-the-Martyr	Holborn	183	66	4	16
Amwell P	Clerkenwell	192	68	11	17
Somer's Town	Paneras	194	60	11	14
Tottenham Court WH	Paneras	196	73	17	22
Old Street	St. Luke	204	60	9	17
St. Mary	Marylebone	211	79	25	20
St. James's Square	St. James, Westminster	212	40	20	5
Whitechapel, North W	Whitechapel	216	36	25	28
Town	Bethnal Green	218	36	24	23
City Road	St. Luke	219	52	8	22
Finsbury	St. Luke	223	43	10	27
Mile End New Town W	Whitechapel	227	36	50	32
Saffron Hill	Holborn	231	40	6	11
Goodman's Fields	Whitechapel	237	28	38	23
Rectory W	Marylebone	238	68	25	23
St. Paul	St. George-in-the-East	242	26	41	20
Borough Road WHP	St. George, Southwark	244	2	167	47
Holywell	Shoreditch	254	36	41	25
All Souls H	Marylebone	258	76	13	16
St. Leonard	Shoreditch	259	41	57	29
Artillery	Whitechapel	271	36	46	22
West London, North WHP	West London	275	36	5	29
St. Botolph W	East London	280	36	19	29
Long Acre	St. Martin-in-the-Fields	287	60	16	22
St. Clement Danes H	Strand	287	36	12	24
Spitalfields	Whitechapel	289	36	40	26
St. Giles, North	St. Giles	291	70	34	27
St. Mary	St. George-in-the-East	291	26	34	29
St. James W	Clerkenwell	291	44	12	24
Cripplegate P	East London	303	44	23	23
St. Giles, South W	St. Giles	317	64	22	31
St. Anne, Soho	Strand	327	64	40	19
Whitecross Street	St. Luke	414	52	16	49
St. Andrew, Eastern W	Holborn	423	50	9	33
Berwick Street	St. James, Westminster	432	65	211	16

NOTE.—The letters placed against the names of the sub-districts denote public institutions within their limits; namely, H, for hospital; L, lunatic asylum; P, prison; W, workhouse.

---

[*The Two Diagrams herewith enclosed should have been inserted in the Scientific Committee's Report on the Cholera-Epidemic of 1854, delivered in August last;—that on the Influence of Elevation at p. 16, and that showing a Comparison of Mortality during the two Epidemics of 1849 and 1854, at p. 24.*]

---






# Cholera and Diarrhoea 1853½-1854.

## LONDON.

### DIAGRAMS.

Representing the Mortality from Cholera and Diarrhoea in Districts at different Elevations in London (I) in the 1½ years 1853½ to 1854 and (II) also the mean Mortality from Cholera in the years 1849 and 1853½-54 at different Elevations.

The Figures in the Diagrams express the Number of Deaths from Cholera to 10,000 Inhabitants living at the Elevations expressed in Feet at the sides of the Diagram




I.

# Cholera — mean of 1849 & 1853½-54.

The Dotted Horizontal Lines represent the Series of Numbers deduced from the Elevation and from the Mortality (145) on the Lowest Level.

Equation of the Curve  $C = Y = \frac{1740}{13 + X}$



(II.)

The Length of the black horizontal line represents the Mortality from Cholera and Diarrhoea at the different Elevations indicated by the height in the base of the Diagram. Thus in Districts 45 Feet above the Thames. The Mortality in London in 1853½-54 from Cholera was 20 in 10,000 Inhabitants from Diarrhoea 20 in 10,000 Inhabitants.







DIAGRAM EXHIBITING BY COLOURED VERTICAL LINE THE DEATHS BY CHOLERA AND BY DIARRHOEA

ON EACH DAY OF THE FIVE MONTHS JUNE — OCTOBER FOR THE YEARS 1849 AND 1854.




The Diagram may be read thus: The Deaths on September 2<sup>nd</sup> from Cholera (blue line) were 459.  
The Deaths from Diarrhoea (yellow) were 36. The Deaths from Cholera and Diarrhoea were 495.











THIRD SERIES—*continued.*TABLE IV.—OF MORTALITY from CHOLERA in the  $1\frac{1}{2}$  years 1853 $\frac{1}{2}$ –1854, and the Number of PERSONS to an ACRE, in the SUB-DISTRICTS of LONDON, arranged in the order of the ELEVATION of the GROUND above Trinity High-water Mark.

SUB-DISTRICTS.	Elevation in Feet above Trinity High- water Mark.	Density.	Mortality by Cholera and Diarrhoea in the $1\frac{1}{2}$ Years 1853 $\frac{1}{2}$ –1854 to 10,000 persons living.	
		Persons to an Acre.	Cholera.	Diarrhoea.
Hampstead W - - - -	350	5	13	11
Sydenham - - - -	188?	2	29	13
Norwood W (L) - - - -	128?	4	27	32
St. John [Marylebone] - - - -	124	55	13	22
Kentish Town - - - -	110	14	9	18
Islington West WHHP - - - -	100	39	13	22
Christchurch [Marylebone] . - - -	92	65	18	27
Islington, East . - - -	88	25	9	21
Regent's Park [Pancras] - - - -	87	75	10	15
Pentonville - - - -	84	89	9	18
Paddington, St. Mary W - - - -	82	21	2	20
St. Mary [Marylebone] . - - -	79	211	25	20
Goswell Street [Clerkenwell] - - - -	78	174	9	14
Stamford Hill [Hackney] - - - -	76	9	17	25
Paddington, St. John H - - - -	76	65	26	15
All Souls [Marylebone] H - - - -	76	258	13	16
Cavendish Square - - - -	73	130	10	10
Tottenham Court [Pancras] wH - - - -	73	196	17	22
Stoke Newington - - - -	72	8	11	12
Streatham (L) - - - -	72	3	19	15
St. George, Bloomsbury - - - -	71	138	12	19
St. Giles, North - - - -	70	291	34	27
Dulwich (L) - - - -	68	1	—	6
Amwell [Clerkenwell] P - - - -	68	192	11	17
Golden Square [St. James, Westminster] W. - - - -	68	166	197	14
Rectory [Marylebone] W - - - -	68	238	25	23
St. George-the-Martyr [Holborn] - - - -	66	183	4	16
Berwick Street [St. James, Westminster]. - - - -	65	432	211	16
Hanover Square - - - -	64	45	12	13
St. Anne, Soho - - - -	64	327	40	19
St. Giles, South W - - - -	64	317	22	31
Camden Town W - - - -	62	123	7	30
Somer's Town - - - -	60	194	11	14
Old Street [St. Luke] - - - -	60	204	9	17
Long Acre - - - -	60	287	16	22
May Fair W - - - -	56	95	12	10
Brixton - - - -	56	10	39	15
West Hackney - - - -	55	40	13	15
Hoxton Old Town - - - -	52	150	14	30
Haggerstone, East - - - -	52	91	15	16
City Road [St. Luke] - - - -	52	219	8	22
Gray's Inn Lane [Pancras] H - - - -	52	171	6	15
Whitecross Street [St. Luke] - - - -	52	414	16	49
Hoxton New Town w - - - -	52	181	8	22
Haggerstone, West W - - - -	52	154	12	25
St. Andrew, Eastern [Holborn] W - - - -	50	423	9	33
St. Mary-le-Strand - - - -	48	173	12	19
City of London, North-west - - - -	44	165	7	8
South Hackney - - - -	44	18	12	31

THIRD SERIES—*continued.*TABLE IV.—Of Mortality from Cholera, &c.—*continued.*

SUB-DISTRICTS.	Elevation in Feet above Trinity High- water Mark.	Density.	Mortality by Cholera and Diarrhoea in the 1½ Years 1853½-1854 to 10,000 persons living.	
		Persons to an Acre.	Cholera.	Diarrhoea.
St. James, [Clerkenwell] W - -	44	291	12	24
City of London, North-east - -	44	139	8	16
Hackney W - - - -	44	12	19	19
Cripplegate [East London] P - -	44	303	23	23
Hackney Road [Bethnal Green] - -	44	170	27	18
Finsbury - - - -	43	223	10	27
Lee - - - -	42	4	21	18
St. Leonard [Shoreditch] - -	41	259	57	29
St. James's Square [St. James, West- minster]. - - - -	40	212	20	5
Saffron Hill [Holborn] - - - -	40	231	6	11
LONDON - - - -	39	30	46	25
Artillery [Whitechapel] - - - -	36	271	46	22
Church [Bethnal Green] - - - -	36	165	16	18
St. Botolph [East London] W - -	36	280	19	29
St. Clement Danes H - - - -	36	287	12	24
Mile End New Town W - - - -	36	227	50	32
Spitalfields - - - -	36	289	40	26
West London, North WwHP - - -	36	275	5	29
Whitechapel, North W - - - -	36	216	25	28
Green [Bethnal Green] W - - - -	36	60	27	30
Town [Bethnal Green] - - - -	36	218	24	23
Holywell [Shoreditch] - - - -	36	254	41	25
Mile End Old Town, Upper - - -	32	155	16	22
Whitechapel Church H - - - -	32	166	21	16
Goodman's Fields - - - -	28	237	38	29
Mile End Old Town, Lower W - -	28	55	26	33
Kensington Town WL - - - -	28	23	49	27
St. Mary [St. George in the East] -	26	291	34	29
St. Paul [St. George in the East] -	26	242	41	20
West London, South - - - -	24	178	12	9
City of London, South-west - - -	21	137	18	10
City of London, South - - - -	21	115	18	9
City of London, South-east - - -	21	103	24	10
Clapham - - - -	21	13	109	21
Aldgate - - - -	19	99	107	34
Ratcliff W - - - -	18	115	64	28
Charing Cross WH - - - -	17	48	26	13
Lewisham Village Ww - - - -	16	2	41	28
Bow w - - - -	15	13	44	26
Chelsea, North-east L - - - -	13	70	43	27
Chelsea, North-west Ww - - - -	12	83	38	24
Belgrave [St. George, Han. Square.] H -	12	69	49	21
Greenwich, West P - - - -	12	58	35	26
Brompton H - - - -	12	21	34	17
Wandsworth P - - - -	12	4	70	34
Putney - - - -	12	2	18	11
Chelsea, South - - - -	10	52	76	24
St. Paul, Deptford - - - -	10	15	45	24
Limehouse W - - - -	10	86	50	19
Hammersmith, St. Paul - - - -	8	6	80	20
Kennington, 2d - - - -	8	37	81	16
Shadwell W - - - -	7	90	69	37



THIRD SERIES—*continued.*TABLE IV.—Of Mortality from Cholera, &c.—*continued.*

[SUB-DISTRICTS.]	Elevation in Feet above Trinity High- water Mark.	Density.  Persons to an Acre.	Mortality by Cholera and Diarrhoea in the 1½ Years 1853½-1854 to 10,000 persons living.	
			Cholera.	Diarrhoea.
Greenwich, East W - - -	7	10	62	54
Fulham w L - - -	6	6	59	25
St. Olave, Southwark H - -	6	107	163	26
Camberwell W - - -	5	13	120	37
Hammersmith, St. Peter - -	4	22	83	12
Peckham - - -	4	17	101	26
St. Nicholas, Deptford - -	4	47	206	46
St. Margaret, Westminster WHP -	4	48	54	33
Kennington, 1st - - -	4	53	135	37
St. Saviour, Southwark H - -	4	127	208	73
Battersea W - - -	3	5	152	49
Poplar W - - -	3	19	42	25
Waterloo Road, 1st - - -	3	155	49	23
St. John [St. George-in-the-East] W -	2	103	33	53
St. John [Westminster] HP - -	2	132	63	28
Lambeth Church, 1st - - -	2	89	39	32
Waterloo Road, 2d - - -	2	129	78	36
Christchurch [Southwark] W - -	2	169	75	29
St. John, Horsleydown W - -	2	121	140	14
Borough Road WP - - -	2	244	167	47
Lambeth Church, 2d W - -	1	114	63	46
London Road [St. George, Southwark] -	0	159	71	21
Rotherhithe W - - -	0	20	171	54
Leather Market [Bermondsey] - -	0	166	179	44
St. Mary Magdalen [Bermondsey] W -	0	98	169	47
St. Mary [Newington] - - -	-1	87	74	11
St. James [Bermondsey] - - -	-1	42	201	29
Trinity [Newington] P - - -	-1	147	115	32
Kent Road - - -	-1	173	142	37
St. Peter, Walworth W - - -	-2	93	136	34
St. George [Camberwell] - - -	-3	37	94	28

NOTE.—The letters placed against names of sub-districts denote public institutions within their limits; *W* stands for workhouse; *H*, for hospital; *L*, lunatic asylum; *P*, prison; *w* indicates a workhouse not belonging to the district, though situated therein.

A minus sign (–) before the figures indicates that the level is *below* Trinity High-water Mark.

The elevation of Eltham, Plumstead, Woolwich Dockyard, and Woolwich Arsenal has not been determined, the mortality from Cholera in 1853½ to 1854 was respectively 12, 16, 12, and 43 to every 10,000 inhabitants, and the mortality from Diarrhoea was respectively 8, 22, 16, and 20.

## THIRD SERIES—continued.

TABLE V.—Showing the MORTALITY by CHOLERA in 1849 and in the 1½ years 1853½–1854 of FOURTEEN SECTIONS of the POPULATION of LONDON living in SUB-DISTRICTS at FOURTEEN different STAGES of ELEVATION; also other CONDITIONS.

Annual Value of Houses.	Elevation in Feet above Trinity High- water Mark of the Thames.		Density.	Annual Rate of Increase per Cent. on Population 1841-51.	Mortality by Cholera and Diarrhœa 1853½-1854.			Mortality by Cholera.	
			Persons to an Acre in 1851.		Deaths to 10,000 Persons.				
					Cho- lera.	Diarr- hœa.	Cholera and Diarr- hœa.	In 1849.	Mean in the two Epidem- ics.
£	Feet.	Feet.							
40	100 —	350	13	5·761	13	21	34	12	13
71	90 —	100	65	1·634	18	27	45	10	14
38	80 —	90	33	4·161	8	19	27	23	16
48	70 —	80	33	*883	19	18	37	25	22
70	60 —	70	79	*901	36	20	56	26	31
36	50 —	60	67	2·607	13	22	35	45	29
53	40 —	50	37	*885	20	20	40	44	32
27	30 —	40	170	1·207	25	25	50	77	51
60	20 —	30	43	1·963	40	23	63	43	44
40	10 —	20	19	2·709	50	24	74	60	55
28	Under 10, viz. :		32	1·958	104	35	139	121	113
31	5 —	10	13	2·040	85	31	116	91	88
32	3 —	5	24	2·182	103	36	139	87	95
31	1 —	3	132	1·326	78	35	113	138	108
21	—3 —	1	64	2·264	137	34	171	153	145
41	Total of London		30	1·982	46	25	71	62	54

TABLE VI.—Showing the MORTALITY by CHOLERA in 1849 and in the 1½ years 1853½–1854 of SIX PORTIONS of the POPULATION of LONDON living in SUB-DISTRICTS at Six different Elevations; and other Conditions.

No. of Sub-Districts.	Elevation above Trinity High-water Mark of the Thames.			Density  Persons to an Acre in 1851.	Annual Value of Houses.	Annual Rate of Increase per Cent. on Population 1841-51.	Mortality by Cholera and Diarrhoea 1853½-1854.			Mortality by Cholera.	
	Extreme Elevation of Sub-districts.		Mean.				Deaths to 10,000 Persons living.			Deaths to 10,000 persons living.	
							Cholera.	Diarrhoea.	Cholera and Diarrhoea.	In 1849.	Mean in the two Epidemics.
6	Feet. 100	Feet. — 350	Feet. 137	13	£ 40	5·761	13	21	34	12	13
5	80	— 100	88	37	45	3·496	10	21	31	20	15
24	60	— 80	69	48	60	*893	28	19	47	25	26
24	—	60	48	47	45	1·689	17	21	38	45	31
23	20	— 40	31	81	41	1·527	32	24	56	65	49
49	Under 20		5	27	32	2·178	88	31	119	103	96
ALL LONDON	{ 3 ft. below to } { 350 feet above }		39	30	41	1·982	46	25	71	62	54
1	2		3		5	6	7	8	9	10	11

The column 2 shows that the mean elevation of the sub-districts ranges between 100 feet and 350 feet; and the 3d column, that upon multiplying the population of each sub-district into its elevation, and dividing all the sums thus obtained by the sum of the population, the mean elevation at which the people lived in the six highest sub-districts is about 137 feet.

THIRD SERIES—*continued.*

TABLE VII.—Of the PRINCIPAL FACTS from which the RESULTS in Tables V. and VI. were obtained.

Elevation above Trinity High-water Mark of the Thames.		Area.	Population.		Deaths from Cholera and Diarrhoea, 1853½–1854.		
			1841.	1851.	Cholera.	Diarrhoea.	Cholera and Diarrhoea.
Feet.	Feet.	Acres.					
100 —	350	9,341	70,119	121,497	194	299	493
90 —	100	518	28,311	33,395	64	98	162
80 —	90	3,287	72,740	108,522	98	239	337
70 —	80	5,919	177,169	193,128	373	360	733
60 —	70	2,844	206,649	225,663	835	455	1,290
50 —	60	2,854	147,501	189,876	273	463	736
40 —	50	5,363	181,400	197,791	410	405	815
30 —	40	1,311	197,680	222,386	573	581	1,154
20 —	30	3,523	140,510	170,051	726	408	1,134
10 —	20	13,637	197,082	256,182	1,385	660	2,045
Under 10, viz.:	*	18,429	492,000	595,119	6,609	2,187	8,796
5 —	10	7,742	83,820	102,191	933	342	1,275
3 —	5	6,693	128,890	159,298	1,764	619	2,383
1 —	3	1,145	132,745	151,070	1,226	558	1,784
—3 —	1	2,849	146,545	182,560	2,686	668	3,354
Elevations not stated	-	67,026	1,911,761	2,314,110	11,540	6,155	17,695
Police on duty	-	11,003	33,506	48,126	121	103	224
	-	—	3,090	—	—	—	—
Total	-	78,029	1,948,417	2,362,236	11,661	6,258	17,919

\* The numbers in this line are the sums of the numbers in the four lines below. The last line (—3—1) is to be read thus: the mean elevations of the lowest sub-districts range from three feet below to one foot above Trinity High-water Mark.

TABLE VIII.—Estimated Number of CASES of CHOLERA and DIARRHOEA in London in the year and half, July 1st, 1853, to the end of 1854; also the Deaths registered.

Ages.	Cholera and Diarrhoea.		Cholera.		Diarrhoea.	
	Cases (estimated).	Deaths (registered).	Cases (estimated).	Deaths (registered).	Cases (estimated).	Deaths (registered).
0—	74,424	6,393	3,465	2,142	70,959	4,251
5—	15,210	1,237	2,178	1,056	13,032	181
10—	6,173	513	968	452	5,205	61
15—	13,356	1,229	3,108	1,086	10,248	143
25—	38,228	1,826	4,624	1,637	33,604	189
35—	74,001	1,979	4,127	1,793	69,874	186
45—	46,687	1,584	2,747	1,376	43,940	208
55—	54,953	1,434	2,103	1,132	52,850	302
65—	13,302	1,079	1,179	686	12,123	393
75—	10,617	546	374	267	10,243	279
85 and upwards	7,744	99	44	34	7,700	65
All ages	354,695	17,919	24,917	11,661	329,778	6,258

THIRD SERIES—*continued.*

TABLE IX.—Showing the PROBABILITY of RECOVERING and of DYING at different PERIODS of CHOLERA (deduced from the Medical Returns).

Time from Commencement.	Remaining in the Period following.	To Recover.	To Die.	Terminating in the period indicated in Column 1.	Recovering.	Dying.
Hours.						
0—	3,600	1,856	1,744	106	31	75
6—	3,494	1,825	1,669	347	32	315
12—	3,147	1,793	1,354	274	17	257
18—	2,873	1,776	1,097	89	2	87
Days.						
0—	3,600	1,856	1,744	816	82	734
1—	2,784	1,774	1,010	590	162	428
2—	2,194	1,612	582	403	217	186
3—	1,791	1,395	396	342	236	106
4—	1,449	1,159	290	271	197	74
5—	1,178	962	216	185	143	42
6—	993	819	174	163	122	41
7—	830	697	133	30	20	10
8—	800	677	123	28	20	8
9—	772	657	115	27	21	6
10—	745	636	109	25	20	5
11—	720	616	104	24	19	5
12—	696	597	99	22	18	4
13—	674	579	95	21	17	4
14—	653	562	91	89	59	30
15—	564	503	61	78	59	19
16—	486	444	42	76	66	10
17—	410	378	32	69	62	7
18—	341	316	25	65	60	5
19—	276	256	20	60	56	4
20—	216	200	16	51	48	3
21—	165	152	13	30	28	2
22—	135	124	11	23	21	2
23—	112	103	9	17	16	1
24—	95	87	8	15	14	1
25—	80	73	7	12	11	1
26—	68	62	6	10	9	1
27—	58	53	5	10	9	1
28—	48	44	4	10	9	1
29—	38	35	3	9	8	1
30—	29	27	2	9	8	1
31—	20	19	1	8	7	1
32—	12	12	0	6	6	0
33—	6	6	0	4	4	0
34—	2	2	0	2	2	0
1	2	3	4	5	6	7

The Table may be read thus :—By the Medical Returns of 3,600 persons that are attacked by Cholera, 1,856 ultimately recover, 1,744 die ; 816 terminate on the first day, 82 by recovery, 734 by death, leaving 2,784 to enter the second day, of whom 1,774 will recover, 1,010 will die, &c.

The cols. 2, 3, 4, it will be observed, are derived from the first numbers at their head by the successive subtraction of the numbers furnished by the returns in cols. 5, 6, 7.



THIRD SERIES—*continued.*

TABLE X.—DAILY RATE of RECOVERY and MORTALITY at different PERIODS of CHOLERA.

Day of the Disease.	To 100 Patients living at each period, the Number of Cases		
	Daily Terminating.	Daily Recovering.	Daily Dying.
Days.			
0 — 1	26	3	23
1 — 2	24	7	17
2 — 3	20	11	9
3 — 4	21	14	7
4 — 5	21	15	6
5 — 6	17	13	4
6 — 7	18	13	5
7 — 8	4	3	1
8 — 9	4	3	1
9 — 10	4	3	1
10 — 11	3	2	1
11 and up <sup>s</sup> .	11	10	1

The Table may be read thus :—To every 100 patients in the *first day* of the disease, 3 recoveries, 23 deaths took place ; to every 100 patients in the *second day* of the disease 7 recoveries, 17 deaths took place. Thus the rate of mortality was highest on the first and second days ; and the rate of recovery was highest on the fifth day (4-5.) If there were 12 hospitals, each containing 100 beds, constantly occupied by patients in the first, second, third, &c. day of the disease, there would be 3 recoveries, 23 deaths in the first, &c.

THIRD SERIES—*continued.*

TABLE XI.—Showing the PROBABILITY of RECOVERING or DYING from an Attack of CHOLERA at different PERIODS of the DISEASE.

Time from Commencement.	Probability of ultimately	
	Recovering.	Dying.
Hours.		
0—	·5156	·4844
6—	·5223	·4777
12—	·5698	·4302
18—	·6182	·3818
Days.		
0—	·5156	·4844
1—	·6372	·3628
2—	·7347	·2653
3—	·7789	·2211
4—	·7999	·2001
5—	·8166	·1834
6—	·8248	·1752
7—	·8398	·1602
8—	·8463	·1537
9—	·8510	·1490
10—	·8537	·1463
11—	·8556	·1444
12—	·8578	·1422
13—	·8591	·1409
14—	·8606	·1394
15—	·8918	·1082
16—	·9136	·0864
17—	·9220	·0780
18—	·9267	·0733
19—	·9275	·0725
20—	·9259	·0741
21—	·9212	·0788

The Table may be read thus :—The average probability that a patient who has lived two complete days (48 hours) from the commencement of the attack of cholera will ultimately recover, is ·7347; or it is 7347 to 2653 in favour of his recovery.

## No. III.

**Summary Table of Houses, Population, and Mortality, in the  
Golden Square Districts.**

*(Prepared by Mr. H. EDWARDS, from the Registrar-General's Returns.)*

SUMMARY TABLE showing the HOUSES, POPULATION, and DEATHS from all Causes, from Zymotic Diseases, and from Cholera especially, in each Street, Court, &c., in the Registrar's Districts of St. Ann's, Soho, Berwick Street, and Golden Square, during the Seven Years 1848-1854, both inclusive.

Name of Street, Court, &c.	Population of Street, 1851.	Total No. of Houses in Street.	Number of HOUSES in which Deaths have occurred during the seven years 1848-1854.				Number of DEATHS during the seven years 1848-1854.				Total Cholera Deaths in the Epidemic of 1853-4.	Number of HOUSES in which Deaths from Cholera occurred in the periods 1848-52, and 1848-9.		
			From Zymotic Diseases.		From Cholera.	Exclusive of Cholera.	From All Causes.	From Zymotic Diseases.	From Cholera alone.	In Houses where other Zymotic Deaths have occurred.	In Houses where no other Zymotic Deaths have occurred.	1848-52.	1853-4.	1848-9 and 1853-4.
			From All Causes.	From Zymotic Diseases.	Exclusive of Cholera.	From Zymotic Diseases.	From Cholera alone.	From Cholera alone.	From Cholera alone.	From Cholera alone.	From Cholera alone.	From Cholera alone.	From Cholera alone.	From Cholera alone.
Aberdeen Mews	16	3	2	-	-	3	2	-	-	-	-	-	-	-
Allen Court	6	1	1	-	-	2	2	-	-	-	-	-	-	-
Archer Street	294	18	14	3	2	48	7	1	1	1	1	1	1	-
Argyle Place	113	11	5	-	-	6	-	-	-	-	-	-	-	-
Arundel Street and Pantons Square.	118	20	9	1	-	10	1	-	-	-	-	-	-	-
Argyle Street	244	34	7	2	-	15	2	-	-	-	-	-	-	-
Bateman's Buildings	200	16	12	2	-	17	2	-	-	-	-	-	-	-
Bear Street	49	5	5	2	1	9	2	-	-	-	-	-	-	-
Bentink Street	327	13	14	4	7	51	11	5	7	1	7	7	7	-
Berwick Street	1,524	102	77	17	29	244	52	25	25	25	25	25	25	-
Blenheim Mews	32	6	3	1	-	4	1	-	-	-	-	-	-	-
Blenheim Street	191	16	12	5	1	21	8	-	-	-	-	-	-	-
Brewer Street	246	26	14	2	1	29	2	-	-	-	-	-	-	-
Bridle Lane	68	35	7	1	3	12	1	-	-	-	-	-	-	-
Broad Street	855	49	45	9	29	190	31	29	31	29	31	29	29	-
Brown's Court	17	1	2	-	-	2	-	-	-	-	-	-	-	-
Burlington Gardens	11	2	2	-	-	3	-	-	-	-	-	-	-	-
Burlington Street	-	-	1	-	-	1	-	-	-	-	-	-	-	-
Burlington Mews	-	-	2	1	-	2	-	-	-	-	-	-	-	-
Cambridge Street	173	15	13	3	6	39	6	-	-	-	-	-	-	-



Carnaby Street	-	965	58	51	17	14	162	33	14	5	17	2	12	-	-
Carlisle Street	-	175	13	11	3	-	22	3	-	-	-	-	-	-	-
Castle Street	-	36	4	4	1	-	4	1	-	-	-	-	-	-	-
Catherine-wheel Yard	-	31	5	4	2	-	10	2	-	-	-	-	1	-	-
Chapel Place	-	177	20	16	5	1	29	8	1	-	1	-	-	-	-
Charles Street	-	90	12	3	-	-	4	-	-	-	-	-	1	-	-
Church Street	-	538	35	28	11	2	74	5	2	-	1	1	1	-	-
Clifford Street	-	114	18	10	11	-	13	-	-	-	-	-	-	-	-
Coach and Horses Yard	-	30	35	3	1	-	3	1	-	-	-	-	-	-	-
Cock Court	-	58	3	3	-	-	6	-	-	-	-	-	-	-	-
Copper's Court, or Pugh's Place	-	42	6	5	1	-	9	1	-	-	-	-	-	-	-
Cork Street	-	202	23	7	3	-	12	-	-	-	-	-	-	-	-
Coventry Street	-	130	20	6	3	-	12	3	-	-	-	-	-	-	-
Cranbourne Passage	-	31	4	2	2	-	2	2	-	-	-	-	-	-	-
Cranbourne Street	-	147	25	11	2	-	13	3	-	-	-	-	-	-	-
Cross Court	-	30	4	3	3	1	4	3	-	1	1	1	1	-	-
Cross Street	-	367	14	13	4	5	71	10	4	6	9	1	5	-	-
Crown Court	-	132	9	12	6	1	37	8	2	-	-	1	1	-	-
Crown Street	-	642	40	30	11	5	111	6	3	2	1	4	-	-	-
Dean Street	-	947	96	63	24	3	123	32	-	3	-	3	-	-	-
Devonshire Place	-	13	1	2	-	-	5	-	-	-	-	-	-	-	-
Down's Place	-	40	9	2	-	-	2	-	-	-	-	-	-	-	-
Duck Lane	-	21	21	3	-	2	4	-	-	2	2	-	2	-	-
Dufour's Place	-	170	10	10	2	3	27	4	2	3	4	1	3	-	-
Earl's Court	-	50	6	3	1	-	3	1	-	-	-	1	-	-	-
Edward Street	-	95	9	7	1	4	15	-	-	4	3	1	3	-	-
Falconberg Court	-	139	9	9	-	6	31	16	6	-	-	6	-	-	-
Faulbert's Place	-	28	4	2	-	-	2	-	-	-	-	-	-	-	-
Frith Street	-	609	65	28	3	-	36	3	-	-	-	-	-	-	-
George Court	-	81	8	6	-	-	12	-	-	1	-	1	1	-	-
George Place (Cross Street)	-	58	4	5	2	1	10	5	1	-	1	-	1	-	-
George Street	-	-	-	3	-	1	3	3	-	1	1	-	1	-	-
George Yard	-	47	8	4	2	-	11	-	-	-	-	-	-	-	-
Gerrard Street	-	513	50	30	9	2	54	10	-	2	1	1	1	-	-
Glasshouse Street	-	26	4	2	-	-	4	-	-	-	-	-	-	-	-
Golden Place	-	10	2	2	-	-	2	-	-	-	-	-	-	-	-
Golden Square H	-	370	46	13	2	-	58	-	-	-	-	-	1	-	-
Grafton Street	-	464	27	22	13	4	80	24	5	-	8	3	1	-	-

Summary Table, showing the Houses, Population, and Deaths from all Causes, &amp;c.—continued.

Name of Street, Court, &c.	Popu- lation of Street, 1851.	Total No. of Houses in Street.	Number of HOUSES in which Deaths have occurred during the seven years 1848-1854.			Number of DEATHS during the seven years 1848-1854.			Total Cholera Deaths in the Epidemic of 1853-4.		Number of Houses in which Deaths from <i>Cholera</i> occurred in the periods 1848-52 and 1853-4.			
			From Zymotic Diseases.		From All Causes.	Exclu- sive of Cho- lera.	From ZYMOTIC DISEASES.		From All Causes.	Exclu- sive of Cho- lera.	From Houses where other Zymotic Deaths have occurred.	From Houses where no other Zymotic Deaths have occurred.	1848-52. 1853-4.	1848-9 and 1853-4.
			From All Causes.	Cholera.			From All Causes.	From <i>Cholera</i> alone.						
Great Burlington Street	-	-	1	-	-	1	-	-	-	-	-	-	-	
Great Chapel Place	-	-	1	-	-	1	-	-	-	-	-	-	-	
Great Chapel Street	337	24	17	4	3	48	6	1	3	2	2	1	-	
Great Crown Court	-	27	17	7	1	39	9	-	2	2	2	2	-	
Great Newport Street	-	12	4	-	2	8	-	-	2	-	1	1	-	
Great Marlborough Street	-	55	32	6	5	52	7	-	9	8	4	4	-	
Great Pulteney Street	-	42	30	6	14	81	10	3	18	19	13	1	-	
Great Windmill Street	-	29	18	7	1	59	8	-	1	1	1	1	-	
Greek Street	-	64	37	14	2	74	16	-	2	1	1	1	-	
Green's Court and Pulteney Place.	-	11	12	3	6	42	12	3	5	5	4	4	-	
Green Dragon Yard	-	2	3	1	-	4	1	-	-	-	-	-	-	
Ham Yard (Duck Lane)	-	3	1	-	-	1	-	-	-	-	-	-	-	
Harris Place	-	4	2	-	-	5	-	-	-	-	-	-	-	
Hayes Court	-	12	6	1	-	10	1	-	-	-	-	3	-	
Heddon Court	-	12	14	6	3	37	9	1	2	3	-	-	-	
Heddon Street	-	18	15	7	-	40	10	-	-	-	-	-	-	
Hollen Street	-	16	15	4	2	61	9	1	1	1	1	1	-	
Hop Garden	-	-	1	-	-	1	-	-	-	-	-	-	-	
Hopkins Street	-	11	8	2	4	37	9	5	-	4	3	3	-	
Horse and Dolphin Yard	-	2	2	1	1	3	1	-	1	1	1	1	-	
Husband Street	-	10	9	2	2	37	5	2	3	5	2	2	-	
John's Court	-	-	1	-	-	1	-	-	-	-	-	-	-	



Summary Table, showing the Houses, Population, and Deaths from all Causes, &amp;c.—continued.

Name of Street, Court, &c.	Popu- lation of Street, 1851.	Total No. of Houses in Street.	Number of Houses in which Deaths have occurred during the seven years 1848-1854.			Number of DEATHS during the seven years 1848-1854.				Total Cholera Deaths in the Epidemic of 1853-4.	Number of Houses in which Deaths from <i>Cholera</i> occurred in the periods 1848-52 and 1853-4.		
			From all Causes.	From Zymotic Diseases.		From all Causes.	From Zymotic Diseases.				1848-52.	1853-4.	1848-9 and 1853-4.
				From all Causes.	Exclu- sive of Cho- lera.		From Zymotic Diseases.	Cholera.					
Nassau Street	72	10	7	-	1	9	-	-	1	1	-	1	
Naylor's Yard	15	3	3	1	1	3	1	-	1	1	-	1	
New Bond Street	73	7	3	1	-	3	-	-	-	-	-	-	
New Burlington Mews	27	4	1	-	-	1	-	-	-	-	-	-	
New Burlington Street	145	25	4	-	-	4	-	-	-	-	-	-	
Newport Court	161	27	7	2	1	9	-	-	1	1	-	1	
Newport Street	-	-	2	2	-	8	6	-	-	-	-	-	
Newport Market	-	-	10	3	-	12	3	-	-	-	-	-	
New Street	340	18	13	5	8	59	15	10	2	12	-	8	
Noel Street	444	27	21	4	3	56	7	1	4	4	1	2	
Old Burlington Mews	64	11	5	3	-	5	-	-	-	-	-	-	
Old Burlington Street	274	33	11	1	-	18	1	-	-	-	-	-	
Old Compton Street	742	49	33	9	-	83	11	-	-	-	-	-	
Orange Court	22	2	1	1	-	5	2	-	4	1	2	-	
Oxford Street (part of)	866	129	59	13	3	81	16	-	4	12	3	7	
Peter Street	640	31	27	4	10	114	17	12	4	12	-	-	
Philips Court	27	3	2	-	-	2	-	-	-	-	-	-	
Plough Yard	21	1	3	1	-	4	1	-	-	-	-	-	
Poland Street	789	63	49	7	12	107	10	4	8	22	12	-	
Porter Street	503	28	23	9	4	88	17	3	11	1	3	1	
Portland Mews	58	15	1	1	1	11	1	4	-	4	1	1	
Portland Street	523	34	26	6	5	80	12	1	6	7	5	-	
Prince's Court	33	4	3	-	-	9	-	-	-	-	-	-	



Prince's Row	-	488	26	18	10	4	69	15	3	3	3	3	1
Prince's Street	-	530	67	29	10	-	67	-	-	-	-	-	1
Pulteney Court	-	153	8	7	5	5	36	3	3	-	-	-	1
Pulteney Street	-	-	-	8	-	1	8	-	-	-	-	-	-
Queen's Head Court	-	94	7	5	-	2	20	4	1	1	2	1	1
Queen's Street	-	428	35	18	7	1	42	7	1	1	1	1	-
Regent Street (part of)	-	1,092	127	43	11	2	53	12	-	2	-	-	-
Rose Court	-	-	-	4	-	-	4	-	-	-	-	-	-
Richmond Buildings	-	170	13	9	4	1	23	5	-	1	-	-	-
Richmond Street	-	231	10	12	3	1	28	5	-	1	-	-	-
Rose Street	-	265	15	12	6	2	43	2	-	1	-	-	-
Rupert Street	-	802	65	43	20	5	99	-	-	4	-	-	-
Ryder's Court	-	173	17	8	4	-	18	8	1	-	-	-	-
St. Ann's Court	-	543	27	27	6	17	123	25	17	9	-	-	1
St. Ann's Place	-	13	2	4	2	1	13	3	2	-	-	-	-
Saville Passage (Saville Row)	-	-	-	1	-	-	1	4	-	-	-	-	-
Saville Row	-	221	29	9	4	-	13	-	-	-	-	-	-
Ship Yard	-	36	2	2	2	-	4	2	-	-	-	-	-
Silver Street	-	587	48	32	10	11	88	19	-	9	-	-	-
Sutton Place	-	28	6	3	2	-	9	2	-	-	-	-	-
Shooting Gallery (Leicester Square)	-	-	-	1	-	-	1	-	-	-	-	-	-
Smith's Court	-	-	-	1	-	-	1	-	-	-	-	-	-
Soho Square	-	255	36	17	4	-	29	5	-	-	-	-	-
South Row	-	178	7	7	-	6	52	9	11	2	-	-	-
Spur Street	-	25	7	1	-	-	1	-	-	-	-	-	-
Star Court	-	45	4	4	2	-	5	2	-	-	-	-	-
Star Street	-	-	-	1	1	-	1	1	-	-	-	-	-
Sutton Street	-	120	8	8	3	-	19	3	-	-	-	-	-
Sydney Alley	-	23	2	2	2	-	2	2	-	-	-	-	-
Swallow Passage	-	7	1	1	-	-	1	-	-	-	-	-	-
Tichfield Street	-	52	5	3	-	-	6	-	-	-	-	-	-
Tyler's Court	-	116	9	5	23	1	18	4	1	-	-	-	-
Tyler's Street	-	136	16	9	3	2	25	6	2	1	-	-	-
Upper John Street	-	106	10	8	2	1	14	2	-	1	-	-	-
Upper James Street	-	105	7	7	3	3	17	3	1	2	-	-	-
Upper Rupert Street	-	193	10	10	7	-	28	8	-	-	-	-	-

Summary Table, showing the Houses, Population, and Deaths from all Causes, &c.—*continued*.

Name of Street, Court, &c.	Popu- lation of Street, 1851.	Total No. of Houses in Street.	Number of Houses in which Deaths have occurred during the seven years 1848-1854,			Number of DEATHS during the seven years 1848-1854,			Total Cholera Deaths in the Epidemic of 1853-4.	Number of Houses in which Deaths from <i>Cholera</i> occurred in the periods 1848-52 and 1853-4.			
			From Zymotic Diseases.		From all Causes.	From all Causes.	From ZYMOTIC DISEASES.			1848-52.	1853-4.	1848-9 and 1853-4.	
			From all Causes.	Exclu- sive of Cho- lera.			Exclu- sive of Cho- lera.	From all Causes.					From <i>Cholera</i> alone.
Vigo Street -	41	6	1	1	2	1	-	-	-	-	-	-	
Walker's Court -	76	12	3	1	4	1	-	-	-	-	-	-	
Wardour Street -	1,383	124	91	33	183	45	6	12	17	1	11	-	
Warwick Street -	301	49	17	5	22	3	-	-	-	-	-	-	
West Street -	496	26	18	6	68	16	4	7	10	1	4	1	
White Hart Street -	-	-	1	-	1	-	-	-	-	-	-	-	
William and Mary Yard -	110	15	7	4	13	4	-	2	2	-	2	-	
Workhouse (Poland Street) -	628	-	-	1	816	83	44	-	34*	-	-	1	
Total -	42,005	3,485	2,204	686	6,580	1,126	318	323	537	86†	310	10	

\* Ten Deaths occurred in the Workhouse in 1849 and 34 in 1853-4.

† In the 86 houses referred to in the last column but two, there occurred during the years mentioned, 104 deaths, of which 97 were in the epidemic of 1848-9; the others sporadically, 5 in 1850, 1 in 1851, and 1 in 1852. Of the deaths recorded in the last column but three, 11 occurred sporadically in the year 1853, the remainder in the great outbreak of 1854. The "H" affixed to "Golden Square" (p. 3), is intended to intimate that the mortality of a Hospital is included in it.

(Signed) H. EDWARDS.

## No. IV.

Pathological Memorandum, circulated among Medical Officers  
of Public Institutions, and other Members of the Medical  
Profession.

---

THE Medical Council, at the President's desire, have had under consideration what unsettled points in the *pathology of cholera* are at once most important for practical purposes, and most capable of receiving further light from accurate clinical and post-mortem inquiries.

In replying to this reference, the Medical Council are especially desirous that they should not seem in any degree to restrict the field of research, or to discourage independent scrutiny and registration even of the most familiar facts. There is no part of the subject but may be better understood through additional study; and the Medical Council would not willingly dispense with any fruits of careful and conscientious investigation.

With this qualification, the Medical Council would suggest that the physicians of our metropolitan hospitals and the junior officers who work under their direction in the registration of hospital experience, as well as other members of the profession who have the medical charge of public institutions, might, with much utility, collect scientific material in relation to the following questions:—

1. *Through what channel* does the exterior cause or poison of cholera first enter or affect the human body? is it through the lungs? or through the stomach and intestines? or otherwise?

2. Has the disease *a period of incubation*? if so, how long? and on what is it contingent?

3. Is there conclusive evidence, affirmative or negative, as to *communication of the disease from person to person*? Has any disproportionate liability to the disease been suffered by those in attendance on the sick, or by those engaged about their dead bodies, or occupied in cleansing their linen? Have cases of the disease occurred where personal infection was impossible? Have solitary cases arisen in large establishments, or been brought thither, without any diarrhœa or cholera ensuing among other inmates? Where choleraic disease has spread in an establishment, shortly after the arrival either of a choleraic patient or of some person from a choleraic locality, has the establishment previously been free from diarrhœa or fever, and unexceptionable in its sanitary arrangements?

4. Does anything indicate a *communication of the disease by provisions* supplied from houses in which cholera exists?

5. Have persons engaged in *particular manufactures* or other employments appeared to enjoy any special exemption from the disease?

6. Has the disease been observed in apparent *dependence on particular articles of diet*? Has any immunity been enjoyed by persons deriving their water-supply from a different source to that generally supplying their district? Has it occurred to persons who have drunk no other water than such as had previously been distilled, boiled, or filtered through charcoal?

7. Does cholera *begin as a morbid process of the gastro-intestinal mucous membrane*? or is this preceded by some state of general poisoning which requires the gastro-intestinal membrane to act as an emunctory? Is the state of collapse determined by this gastro-intestinal flux, and in proportion to it? or can it arise independently of any such flux? How are the lividity and the cramps determined and proportioned?

8. What conditions determine the occurrence, duration, and severity of *consecutive fever*? What are the varieties of morbid condition included under this term? To what extent does it depend on the previous occurrence of profuse discharges, or on the completeness of collapse? Does stupor in this stage always depend on uræmia, or on what? In what proportion of cases, and under what pathological conditions, is the fever accompanied by exanthem?

9. When *diarrhœa and cholera* prevail together epidemically in a district, are they (with differences of degree) *the same disease*? does the diarrhœa, if left to itself, generally and safely tend to spontaneous recovery? or do such cases, without medical treatment, frequently, in proportion to their numbers, pass into true cholera? Is there any way to discriminate a premonitory diarrhœa.

10. What changes—physical and chemical—are undergone by *the blood in cholera*? Does the consecutive fever represent, in regard to the blood, a period in which this fluid is tending to recover from injuries inflicted on its constitution during the stage of gastro-intestinal flux? Or is it attended by any process of change in the blood, leading to critical discharges or inflammations?

11. Does any *obstruction of the capillary circulation* in the Malpighian tufts of the kidney, or in the lung or brain, or elsewhere, arise either from inspissation or other physical affection of the blood in cholera? Do any infiltrations or other parenchymatous changes, which have been observed in persons dead



from cholera, arise in consequence of such obstructions? Or do all these structural lesions arise as ordinary inflammatory processes?

12. Does the *non-discharge of bile* with the rice-water secretions of cholera depend on tumefaction of the ductus choledochus? or on what?

13. Is the *rise of temperature* which has occasionally been observed *after death by cholera* confined to cases where death occurs during collapse? Does it occur only at the surface of the body, and, if so, depend on a return of blood to the surface? Or does it occur also in the visceral cavities of the body, and in the substance of solid organs?

---









GLASGOW  
UNIVERSITY  
LIBRARY

